

## CAROTENOID-PROTEIN COMPLEXES

P. F. ZAGALSKY

Department of Biochemistry, Bedford College, Regent's Park, London, NW1 4NS, UK

**Abstract**—A survey of carotenoid-protein complexes is presented. Distinction is made between astaxanthin-proteins in which the absorption band of the carotenoid is unaltered in shape, and carotenoid-lipoglycoprotein complexes. The proposal by Buchwald and Jencks<sup>79</sup> that the carotenoid is twisted about the double bonds of the polyene chain, is discussed and sustained for astaxanthin-proteins. A mode of binding for the carotenoid in these complexes is proposed, involving hydrogen-bonding of keto groups of the polyene to imide groups of peptide bonds of the apoprotein.

### INTRODUCTION

This article, following earlier reviews,<sup>1,2</sup> will deal primarily with carotenoid-protein complexes in invertebrates. While they occur elsewhere,<sup>3,4,43,44</sup> it is in the invertebrate phyla that they are widespread. A brief account of "vitamin A-proteins", and plant and bacterial carotenoid-protein complexes is included for completeness.

In invertebrates, carotenoids occur in several different states. They may exist as lipid dispersions, often within chromatophores.<sup>5-7</sup> Vivid and varied coloration may be provided by carotenoids bound to structural elements as, for example, to chitin in the carapace of Crustacea,<sup>8,9</sup> to the scleroproteins of feathers<sup>10,11</sup> and to calcium carbonate in corals.<sup>12,13</sup> Carotenoids are also found as water-soluble complexes with protein. Carotenoid-protein complexes are particularly frequent in the Crustacea, and it is to the complexes of this group that attention has been directed and from which most examples are taken. The taxonomic distribution and anatomical location of carotenoid-protein complexes within animals of the neuston<sup>14</sup> and decapod Crustacea<sup>9</sup> and their possible functions, such as cryptic coloration, within these groups and within the zooplankton,<sup>15</sup> have been amply reviewed.

Two quite different types of carotenoid-protein complex may be distinguished: those in which the carotenoid is associated with a lipo(glyco)protein and those where it is bound stoichiometrically to a simple protein or glycoprotein.

### CAROTENOID-LIPO(GLYCO)PROTEINS

#### Nature of the combinations

The specificity shown by the lipoprotein for the carotenoid component and the nature of the binding, reflected in the spectral characteristics of the complexes, show great variety. In some combinations, such as those in eggs and ovary of *Cancer pagurus* ( $\lambda_{\max}$  470, 490 nm)<sup>16</sup>, in eggs, ovary and blood of *Emerita analoga* ( $\lambda_{\max}$  463, 472 nm)<sup>17</sup> and in eggs of *Pagurus prideauxi* ( $\lambda_{\max}$  470, 495 nm)<sup>156</sup>, selectivity in binding is not apparent. All the carotenoids of the whole tissue are present and in about the same relative proportions. In other instances, as in the ovary of *Pecten maximus* ( $\lambda_{\max}$  472, 498 nm)<sup>16</sup> and plasma of the birds *Ajaia ajaja* ( $\lambda_{\max}$  480 nm)<sup>18</sup> and *Guara*

*rubra* ( $\lambda_{\max}$  460 nm)<sup>11</sup>, polar carotenoids are preferentially combined. In these cases, the bathochromic shift in the absorption spectra (10–25 nm compared with solutions in hexane) can be explained by refractive index effects associated with dissolution of the carotenoids in the lipid or protein components of the lipoproteins.<sup>18,79,82</sup> Part of the carotenoid fraction may nevertheless be associated more specifically. It is of interest that for the carotenoid-transporting low-density lipoprotein<sup>19</sup> of human serum, the circular dichroism induced in the absorption of the carotenoid on cooling has been attributed to its interactions with the protein or to phase transition in the lipid.<sup>19</sup>

A single carotenoid prosthetic group, which may be astaxanthin (*Homarus gammarus* (L.),<sup>1</sup> *Plesionika edwardsi*,<sup>16</sup> *Acanthephyra* spp.,<sup>20</sup> *Giganto cypris*,<sup>21</sup> *Lepas* spp.<sup>1,22,31</sup> and *Pollicipes polymerus*<sup>23</sup>), canthaxanthin (*Artemia salina*<sup>24</sup> and *Branchipus stagnalis* (L)<sup>25</sup>), an epiphasic astaxanthin ester (*Eu*)*Pagurus bernhardus*<sup>26</sup>,†  $\beta$ -carotene (*Callinectes sapidus*)<sup>60</sup> or a "β-carotene-like" carotenoid (*Procambarus* sp.)<sup>27</sup>, is found in many red, green and blue egg and ovary carotenoid-protein complexes. Traces of other carotenoids, however, are often also present<sup>20,23,28,60</sup> and it is debatable whether the apparent specificity is due to the predominance of a single carotenoid in the tissue (*H. gammarus*,<sup>28</sup> *Lepas* spp.,<sup>29</sup> *Acanthephyra* spp.<sup>20</sup> and *C. sapidus*<sup>60</sup>) as a result of metabolic selectivity,<sup>20</sup> or to preferential selection of a particular carotenoid during transport and deposition. The absorption spectrum of the carotenoid in the complexes may have the same spectral characteristics (oscillator strength, half-band width) as the carotenoid dissolved in hexane, with only a small change in the position of the absorption maximum (*A. salina*,  $\lambda_{\max}$  470 nm<sup>24</sup>; *Acanthephyra* spp.<sup>20</sup>,  $\lambda_{\max}$  486–498 nm; *P. polymerus* sp.,  $\lambda_{\max}$  470 nm<sup>23</sup>). Occasionally the smooth absorption band of the prosthetic group may become a triple-peaked band (e.g. in the red complex of *Giganto cypris*<sup>21</sup>) as for ovorubin, the proteinase-inhibiting<sup>30</sup> astaxanthin-glycoprotein of the eggs of the prosobranch *Pomacea canaliculata*.<sup>1</sup> Blue complexes, where the absorption maximum of the keto-carotenoid is bathochromically shifted and altered in shape (usually double-peaked) are common (*P. edwardsi*,  $\lambda_{\max}$  632, (670) nm;<sup>16</sup> *Lepas* spp.  $\lambda_{\max}$  600 nm;<sup>1</sup> *B. stagnalis*  $\lambda_{\max}$  625, (675) nm<sup>25</sup>), while in the spectrum of the green complex of *H. gammarus* (and *Homarus americanus*<sup>41</sup>), ovoverdin  $\lambda_{\max}$  463–468, 645–670 nm nm),<sup>1,28,41</sup> and the purple complex of *P. bernhardus* ( $\lambda_{\max}$  465, 495, 580 nm)<sup>26</sup> absorption bands, altered in shape, occur both above and below (or close to) the band of the free carotenoid.

†The carotenoid-binding specificity of the egg lipoprotein of *P. bernhardus* apparently differs from that of the ovary lipoprotein. The egg carotenoid-lipoprotein complexes of both *P. bernhardus* and *P. prideauxi* have astaxanthin as the major carotenoid component. Substantial amounts of  $\beta$ -carotene and, possibly, traces of astaxanthin esters are also present in the complexes.<sup>156</sup>

The green carotenoid-lipoglycoproteins in anostracan blood (*Branchipus stagnalis* (L.), *Branchinecta packardii* (Pearse), *Tanymastix lacunae* (Guérin) and *Chirocephalus diaphanus* (Prévost))<sup>33</sup> and isopod cuticle (*Idotea montereyensis*<sup>34</sup> and *Idotea resecata*<sup>35</sup>) and epidermis (*Idotea granulosa*<sup>36</sup>) owe their colour to incorporation into the complexes of the complete mixture of tissue carotenoids. The absorption spectra show maxima in the 370, 400–500 and 680 nm regions. In these, it is probable that different modes of binding of the carotenoid exist, for mild treatment, which removes some lipid components, eliminates selectively the 400–500 nm absorption band. It is suggested that lipid-associated carotenoids are responsible for the 400–500 nm absorption, while the maxima in the 370 and 680 nm regions arise from more firmly bound canthaxanthin.<sup>33–36</sup> These absorption bands have different spectral characteristics (half-band width) to those of the carotenoid dissolved in hexane. The green pigment of the carapace of *Carcinus maenas* ( $\lambda_{\max}$  460, 625) similarly show absorption bands above and below that of the free prosthetic group, astaxanthin;<sup>1</sup> the composition of the complex has not been reported. Different modes of binding of carotenoid occur also in the egg and ovary complexes of *P. bernhardus*<sup>26</sup> and *Cyclops vernalis*.<sup>37</sup>

The role of the carotenoid and its mode of binding in these complexes are uncertain. It has been pointed out<sup>17,37–41</sup> that the amount of carotenoid in the lipoproteins, and possibly even the nature of the carotenoid,<sup>40</sup> depends on diet and environmental factors, which influence the availability of carotenoid at vitellogenesis; stoichiometry of combination has seldom been verified and no deleterious effects of carotenoid depletion have been substantiated.<sup>39</sup> Although hatchability<sup>42</sup> of eggs and viability of nauplii<sup>32</sup> may be impaired in animals fed on synthetic diets, this cannot be attributed solely to lack of carotenoids. If definite sites for the carotenoid exist in the proteins, these, as for the vertebrate carotenoid-lipoproteins,<sup>43,44</sup> need not be filled.<sup>41</sup> The approximate stoichiometry in ovooverdin<sup>1,28</sup> and in the egg carotenoid-lipoprotein complexes of *Lepas* sp.<sup>31</sup> and *B. stagnalis*<sup>25</sup> may be fortuitous, and the alteration in the spectrum of the associated keto-carotenoid in these and similar complexes could merely reflect the high polarisability of the polyene.<sup>41</sup> However, the selectivity of association in some of the proteins and the effects of disruption and reconstitution of the carotenoid-lipoprotein linkage (for the complexes of *H. gammarus*, *Lepas* spp. and *P. bernhardus*)<sup>1</sup> suggests that the interaction involves more than passive solution in the lipoprotein.

The instability of the carotenoid-lipoprotein linkage is well documented.<sup>1</sup> Exposure of the proteins to intense light, heat, metal ions or even storage can result in alterations in the visible absorption spectra. Ionic conditions are sometimes found important for the stability of the carotenoid linkage; solutions of the blue lipoprotein complex of *B. stagnalis* immediately turn red unless chloride ions are present.<sup>25</sup> Slight alterations of pH may also affect the spectral characteristics;<sup>41</sup> solutions of the proteins of *Procambarus* spp. turn from brown to orange, reversibly, on changing the pH from 7 to 8<sup>27</sup>, the purple *Sesarma reticulatum* pigment becomes orange at pH 7.5<sup>41</sup> and the purple complex of *P. bernhardus* rapidly loses its

580 nm absorption band even at pH 7.<sup>26</sup> Hypochromicity effects, which may be explained by twisting of the polyene about the single bonds,<sup>109</sup> have been reported in some complexes.<sup>41</sup> The extinction of the carotenoid in the egg lipoglycoprotein of *P. prideauxi* is, however, almost unaltered in the combination.<sup>51</sup>

#### *Lipovitellins and haemolymph lipoglycoproteins*

The high-density lipoglycoproteins, with which most of the carotenoid present in crustacean eggs and ovary may be associated,<sup>1,20,29</sup> have been termed crustacean lipovitellins, in analogy with vertebrate lipovitellins.<sup>41</sup> Purification of the lipoproteins by density gradient and other methods is preferred,<sup>24,27,41</sup> ion-exchange procedures, which can result in removal of lipid and carotenoid,<sup>23,33–36</sup> are to be avoided unless special precautions are taken.<sup>45</sup> The proteins "age" on storage even in the unextracted state.<sup>20,41,47</sup>

The compositions of several crustacean lipovitellins have been investigated and points of similarity revealed.<sup>16,20,27,41,60</sup> They contain 27–35% lipid, a small carbohydrate component and, in contrast to vertebrate lipovitellins,<sup>43</sup> no protein-bound phosphorus.† In most cases, the lipid component consists largely of phospholipid<sup>16,27</sup> with unusual fatty acid composition<sup>27</sup> and smaller quantities of cholesterol and triglycerides,<sup>16</sup> although neutral lipid and phospholipid occur in equal proportion in lipovitellins of *Acanthephyra* spp.<sup>20</sup> The decapod lipovitellins are of similar size, fairly symmetrical in shape and of molecular weight  $3.3\text{--}3.7 \times 10^5$ .<sup>27,41</sup> Lipovitellins of other invertebrates (*P. maximus*;<sup>16</sup> *B. stagnalis*<sup>25</sup>) may be larger or more asymmetrical. Dissociation—reassociation phenomena are observed at low protein concentration.<sup>41</sup> On treatment with sodium dodecylsulphate, more extensive degradation of the lipovitellins of *Procambarus* sp.,<sup>27</sup> *Lepas* sp.,<sup>51</sup> *H. gammarus*,<sup>51</sup> *P. maximus*<sup>51</sup> and *B. stagnalis*<sup>25</sup> is obtained than of amphibian,<sup>48</sup> avian<sup>49</sup> or insect<sup>20</sup> lipovitellins. It should be noted that the electrophoretic patterns obtained depend on the exact conditions of incubation;<sup>51</sup> divergent results have been reported for other lipoproteins.<sup>52</sup>

The amino acid compositions of crustacean vitellins show no unusual features other than high contents of helix-breaking<sup>118,119</sup> amino acids (serine and proline) (Fig. 1). They appear to be homologous proteins with similar contents of non-polar, polar and other categories of amino acids.<sup>56</sup> The compositions are typical of other lipovitellins and soluble lipoproteins (Table 1). An estimate of relatedness can be made by calculating the values of  $S\Delta\phi$  (differences in mol% content of each amino acid squared and summed) for pairs of the proteins.<sup>57,58</sup> Unrelated proteins have  $S\Delta\phi$  values of more than 50 and almost invariably more than 100.  $S\Delta\phi$  values for pairs of vitellins indicate that the crustacean proteins are closely related to each other and to analogous vitellins (Table 2). Their apparent resemblance to the carotenoid-transporting low-density lipoprotein<sup>4</sup> of human serum (LDL), and their distinction from the high-density serum lipoprotein (HDL<sub>s</sub>), may provide an insight into their structure. The protein component of LDL<sub>s</sub> is thought to interact specifically with the cholesteryl moiety of cholesterol esters.<sup>154</sup> In contrast, in the structure of HDL<sub>s</sub>, the protein is associated, in its amphipathic helical regions, with the fatty acid chains of phospholipids and has no intrinsic affinity for cholesterol esters.<sup>155</sup> It is of interest that the carotenoid content of lipoprotein fractions of *Ajaja ajaja* plasma follows that of cholesterol.<sup>18</sup> The lipovitellins are

†The presence of protein-bound phosphorus previously reported,<sup>16</sup> was the result of incomplete lipid extractions of the freeze-dried samples; the phospholipid and total lipid contents quoted in the paper should be increased accordingly.




Fig. 1. Histogram showing the contents (mol %) of amino acids of crustacean lipovitellins: (1) *Homarus gammarus* (L.);<sup>56</sup> (2) *Cancer pagurus*;<sup>56</sup> (3) *Plesionika edwardsi*;<sup>56</sup> (4) *Procambarus* sp.;<sup>27</sup> (5) *Pagurus bernhardus*;<sup>51</sup> (6) *Pagurus prideauxi*;<sup>51</sup> (7) *Lepas* sp.;<sup>31</sup> (8) *Branchipus stagnalis* (L.).<sup>25</sup> The contents of amino acids of the 'average protein'<sup>93</sup> with their standard deviation of occurrence are given by the dashed and dotted horizontal lines, respectively.

Table 1. Comparison of compositions (mol %) of lipovitellins, and some other proteins, in terms of amino acid sets

Proteins	Amino acid sets		P/A ratio	Small§ (mol %)	Charged¶ (mol %)
	Apolar residues (A)† (mol %)	Polarity index (P)‡ (mol %)			
Crustacean lipovitellin					
<i>Homarus gammarus</i>	25.1	52.3	2.1	13.9	33.9
<i>Cancer pagurus</i>	23.6	52.0	2.2	13.9	33.0
<i>Plesionika edwardsi</i>	25.2	54.5	2.2	13.9	34.3
<i>Procambarus</i> sp.	24.1	49.8	2.1	12.8	32.1
<i>Lepas</i> sp.	26.6	52.1	2.0	10.1	32.3
<i>Pagurus bernhardus</i>	30.0	47.2	1.6	12.8	31.3
<i>Pagurus prideauxi</i>	27.3	49.8	1.8	13.1	33.5
<i>Branchipus stagnalis</i>	26.9	48.8	1.8	13.9	32.2
Lamellibranch lipovitellin <sup>56</sup>	21.8	51.1	2.3	16.9	31.1
Insect lipovitellin <sup>50</sup>	26.6	55.1	2.1	8.5	38.8
Amphibian lipovitellin <sup>53</sup>	29.5	47.4	1.6	13.4	31.4
Avian lipovitellin ( $\alpha$ ) <sup>53</sup>	28.4	45.2	1.6	13.2	31.1
Ovotubin, <i>Pomacea canaliculata</i> <sup>56</sup>	28.9	49.5	1.7	11.8	34.3
Serum low density (LDL) lipoprotein <sup>¶</sup> (human)	28.3	51.4	1.8	11.6	33.8
Serum high density (HDL) lipoprotein <sup>††</sup> (human)					
	24.7	53.7	2.2	12.4	40.5
Lipoxygenase (pea, <i>Pisum sativum</i> L.)	26.2	49.5	1.9	12.5	33.6

†Apolar residues:<sup>95</sup> Val, Ileu, Leu, Phe, Met.

‡Polarity index:<sup>94</sup> Sum of mol% values of Asp, Thr, Ser, Glu, Lys, His, Arg.

§Small amino acids:<sup>95</sup> Ala, Gly

¶Charged amino acids:<sup>95</sup> Asp, Glu, Arg, Lys.

<sup>¶</sup>Average amino acid composition derived from the values given in Refs. 52 and 55.

<sup>††</sup>Average amino acid composition derived from the values given in Ref. 54.

Table 2. Estimation of relatedness of egg storage proteins (lipovitellins), serum lipoproteins and other proteins (SA $\phi$  values are given in this table). Possibly related proteins are in italic>

	Lipovitellins	<i>H. gammarus</i>	<i>C. pagurus</i>	<i>P. edwardsi</i>	<i>Procambarus</i> sp.	<i>P. prideauxi</i>	<i>P. bernhardus</i>	<i>Lepas</i> sp.	<i>B. stagnalis</i>	Lamellibranch	Amphibian	Avian ( $\alpha$ )	Insect	Other proteins	Ovourubin	Lipoxygenase	LDL	HDL
Crustacean lipovitellin																		
<i>H. gammarus</i>	0	16	20	59	20	24	18	44	36	40	36	70	27	55	43	112		
<i>C. pagurus</i>	16	0	56	94	49	49	29	66	65	61	63	88	50	68	48	115		
<i>P. edwardsi</i>	20	56	0	47	37	40	33	45	26	51	49	90	45	71	78	155		
<i>Procambarus</i> sp.	59	94	47	0	64	39	54	29	68	19	24	77	51	53	55	169		
<i>P. prideauxi</i>	20	49	37	64	0	10	29	51	72	36	31	84	29	66	51	104		
<i>P. bernhardus</i>	24	49	40	39	10	0	27	35	66	20	16	77	25	47	38	121		
<i>Lepas</i> sp.	18	29	33	62	29	27	0	54	57	44	39	79	42	75	60	135		
<i>B. stagnalis</i>	44	66	45	30	51	35	62	0	60	28	32	107	36	57	51	140		
Lamellibranch lipovitellin	36	65	26	68	72	66	57	60	0	40	40	118	62	74	102	127		
Amphibian lipovitellin	40	61	51	19	36	20	44	28	40	0	6	71	29	47	44	120		
Avian lipovitellin ( $\alpha$ )	36	63	49	24	31	16	39	32	40	6	0	75	32	46	50	116		
Insect lipovitellin	70	88	90	77	84	77	79	107	118	71	75	0	33	48	55	108		
Ovourubin, <i>P. canaliculata</i>	27	50	45	51	29	25	42	36	62	29	32	33	0	23	31	125		
Lipoxygenase (pea, <i>Pisum sativum</i> L.)	55	68	71	53	66	47	75	57	74	47	46	48	23	0	30	127		
Serum low density (LDL) lipoprotein (human)	43	48	78	55	51	38	60	51	102	44	50	55	31	30	0	88		
Serum high density (HDL) lipoprotein (human)	112	115	155	169	104	121	135	140	127	120	116	108	125	127	88	0		

also closely related in amino acid composition to the storage glycoprotein, ovourubin<sup>56</sup> (Table 2).

The carotenoid-lipoglycoproteins (female-specific proteins) that accumulate in the blood of certain Crustacea during vitellogenesis have been shown, in some instances, to be electrophoretically identical with (*Paratellus hydrodomous*<sup>59</sup>), related to (*E. analoga*<sup>17</sup>), or both electrophoretically and immunologically indistinguishable from (*Callinectes sapidus*<sup>60</sup>, *Orchestia gammarella*<sup>61</sup>, *Procambarus* spp.<sup>27</sup>) the lipovitellin deposited in the ovary and eggs.

The synthesis and deposition of crustacean lipovitellin is under endocrine control, but the site of synthesis, probably external to the oocyte in some species, is still controversial.<sup>27,41,59-61</sup> The breakdown of lipovitellin during embryonic development has been followed electrophoretically and immunologically.<sup>46,61</sup> The protein and lipid are utilised to satisfy growth and energy requirements<sup>1,20,61</sup> while the carotenoid becomes concentrated in the naupliar eye and chromatophores.<sup>6,9,15,20,29,62</sup> Opinions differ as to whether the carotenoid content of the egg changes during development,<sup>17,20,39,46</sup> although it may be altered chemically.<sup>20</sup>

The lipovitellin is either dispersed in the oocytes of Crustacea (e.g. within spheroid micelles, *Lepas* sp.<sup>29</sup>) or in granular yolk bodies.<sup>63</sup> In eggs of the anostracans, *A. salina*<sup>24</sup> and *B. stagnalis*,<sup>25</sup> however, the canthaxanthin-lipovitellin complex is packaged into small, oval platelets together with large amounts of diguanosine nucleotides and some deoxyribonucleic acid. The nucleotides are utilised for purine synthesis during development.<sup>64,65</sup>

In some Crustacea, carotenoid-lipoglycoproteins (e.g. the green complexes of certain anostracans<sup>35</sup>) occur in the blood of males and females. These pigments, not yet fully characterised, may have a function in colour adaptation.<sup>66</sup> However, purple and green carotenoid-lipoprotein complexes are seen in the blood of *Tanymastix lacunae* in separate individuals (male or female) at the same stage of

growth and in the same culture medium.<sup>32</sup> A function other than that of protective coloration may therefore be fulfilled by these complexes. In another species (*Branchinecta packardi*) carotenoid is mainly present, in the female only, as blue needle-like deposits (carotenoid-protein or lipoprotein complex?).<sup>32</sup>

#### Cuticular, epidermal and other carotenoid-lipoglycoproteins

The green cuticular and epidermal pigments of several idoteid isopods have been identified as lipoglycoproteins containing, in addition to canthaxanthin, several lipid-associated carotenoids.<sup>34-36</sup> While detailed chemical analyses have not been attempted, the protective coloration function of these pigments has been convincingly presented.<sup>35,67</sup>

The chemical composition of the blue epidermal pigment of *Palaemon serratus*, the synthesis of which is under hormonal control,<sup>68</sup> has not been reported.

The blue muscle pigment of the sea squirt, *Salpa cylindrica*, is unusual in having a single-banded absorption spectrum derived from a carotenoid with a typical triple-peaked spectrum.<sup>21</sup> Composition of the protein has not been determined; the unidentified carotenoid fails to combine with the apoproteins of several simple carotenoproteins (Table 5).

#### Enzymes

Enzyme utilising carotenoids as substrates must necessarily occur as "carotenoid-protein complexes".<sup>2,69</sup> With the exception of lipoxygenase of pea (*Pisum sativum* L.),<sup>70</sup> compositions of carotenoid-metabolising enzymes have yet to be reported. The lipoxygenase which oxidises  $\beta$ -carotene in the presence of linoleic acid and O<sub>2</sub>, has an amino acid composition consistent with that of soluble lipoproteins (Tables 1 and 2), with which it may be provisionally classed.

## CAROTENOPROTEINS

There is a clear distinction between the carotenoid-lipoprotein complexes and those associations in which the carotenoid is bound stoichiometrically to a simple protein or, in a single known instance (ovorubin), to a glycoprotein.<sup>1</sup> True carotenoproteins, typified by ovorubin and crustacyanin, the blue pigment of lobster carapace, are stable combinations where the carotenoid tends to stabilise the tertiary and/or quaternary protein structures, a property well known for other polyenes in combination with protein.<sup>87,89</sup> Native carotenoprotein may often be reconstituted from carotenoid and freshly prepared apoprotein.<sup>1</sup>

*Physical properties and subunit structure*

Simple carotenoproteins have been isolated from several invertebrate species and a variety of tissues (Fig. 2). The prosthetic group has been identified invariably as astaxanthin. The position of the absorption maxima of the blue/purple combinations range from 520 nm (carapace carotenoprotein of *Pachygrapsus marmoratus*)<sup>71</sup> to 655 nm (carapace pigment of *Cambarus clarkii*).<sup>75</sup>

Molecular weights of carotenoproteins (native or  $\alpha$ -forms) range from  $5 \times 10^4$  (carapace carotenoprotein of *Aristeus antennatus*) to  $1 \times 10^6$  (asymmetrical mandible carotenoprotein of *A. antennatus*).<sup>71</sup> At low ionic strength many of the proteins dissociate to give purple derivatives,  $\beta$ -forms, of molecular weight  $4.0\text{--}5.0 \times 10^4$ . The apoproteins, prepared simply by acetone precipitation, are often electrophoretically heterogeneous. Their molecular weights, lying between  $1.8$  and  $2.6 \times 10^4$  (crustacyanin;<sup>77-81</sup> mantle carotenoprotein of *Velella* sp.,<sup>31</sup> *A. antennatus*

carotenoproteins;<sup>71</sup> hypodermal carotenoproteins of *Labidocera acutifrons*<sup>72</sup> and *Anomalocera patersoni*<sup>31</sup>) or between  $0.8$  and  $1.0 \times 10^4$  (carapace carotenoprotein of *C. clarkii*<sup>73</sup> and hypodermal carotenoprotein of *Labidocera* sp.<sup>31</sup>), indicate the influence of the prosthetic group on quaternary structure. Each apoprotein may bind one (crustacyanin;<sup>78,79,88</sup> ovorubin;<sup>51</sup> *A. antennatus* carotenoproteins<sup>71</sup>), two (*Velella* sp. and *Anomalocera patersoni* carotenoproteins<sup>31</sup>) or, possibly, three (*Labidocera acutifrons* carotenoprotein<sup>72</sup>) molecules of astaxanthin. Exceptionally, the apoprotein of ovorubin retains the quaternary structure of the carotenoprotein.<sup>51</sup>

A yellow complex isolated from lobster carapace<sup>79-81</sup> has a low minimum molecular weight (based on astaxanthin content) of  $4.4 \times 10^3$  and an absorption maximum at 409 nm, well below that of carotenoid in hexane.<sup>79</sup> The characteristics of the absorption spectrum (half-band width, oscillator strength and extinction coefficient) are quite different from those of the blue/purple carotenoproteins. It has similar absorption and optical rotatory dispersion (ORD) spectra to those of astaxanthin aggregates in which the molecules are believed to be stacked with the dipoles of successive molecules at an angle to one another.<sup>79,82</sup> It is in doubt whether this carotenoid aggregate, kept in solution by a protein component immunologically related to crustacyanin,<sup>79,82</sup> occurs naturally in the carapace.

The five apoprotein units of crustacyanin have been obtained in homogeneous state by ion-exchange chromatography.<sup>83</sup> The two components which migrate the furthest anodically in starch gel electrophoresis at pH 8.6, show similar sizes ( $1.8\text{--}2.0 \times 10^4$ ) in gel filtration<sup>83</sup> and




Fig. 2. Histogram showing the contents (mol %) of amino acids of carotenoproteins: (1) *Cambarus clarkii* carapace;<sup>73</sup> (2) *Labidocera* sp., hypodermis;<sup>31</sup> (3) *Anomalocera patersoni*, hypodermis;<sup>31</sup> (4) *Porpita* sp., mantle and tentacles;<sup>31</sup> (5) *Labidocera acutifrons*, hypodermis;<sup>72</sup> (6) *Velella* sp., mantle;<sup>31</sup> (7) *Homarus gammarus* (L.), carapace, average value;<sup>71,78,79</sup> (8) *Clibanarius erythropus*, exoskeleton;<sup>71</sup> (9) *Scyllarus arctus*, carapace;<sup>71</sup> (10) *Aristeus antennatus*, mandibles;<sup>71</sup> (11) *Aristeus antennatus*, carapace;<sup>71</sup> (12) *Aristeus antennatus*, stomach;<sup>71</sup> (13) *Eriphia spinifrons*, carapace.<sup>71</sup> The content of amino acids of the 'average protein'<sup>93</sup> with their standard deviation of occurrence are given by the dashed and dotted horizontal lines, respectively.

acrylamide-sodium dodecylsulphate electrophoresis.<sup>51</sup> The amino acid compositions differ by only 4  $S\Delta\phi$  units,<sup>83</sup> a value obtainable for identical proteins.<sup>57</sup> The three remaining apoprotein units, two migrating cathodically and one anodically, are also similar to each other in size ( $2.1\text{--}2.3 \times 10^4$ ) and amino acid composition, pairs giving  $S\Delta\phi$  values of 8, 9 and 23.<sup>83</sup>

The two sets of subunits are themselves related in composition, for the major components of each set differ by only 46  $S\Delta\phi$  units, most of which is due to differences in leucine and threonine.<sup>83</sup> The amino acid compositions of  $\alpha$  and  $\gamma$ -crustacyanin are consistent with an equal contribution of each set of subunits to the structures.<sup>83</sup> The latter protein has a higher sedimentation and diffusion constant than  $\alpha$ -crustacyanin and elutes before it in gel filtration. It differs in shape and absorption maximum from  $\alpha$ -crustacyanin, but has an identical amino acid composition, molecular weight and subunit structure.<sup>79,83</sup> All five apoproteins retain the ability to recombine with astaxanthin, singly or in pairs, to form the non-interconvertible dimeric  $\beta$ -crustacyanins. The preferred association is between subunits of the two separate types; these forms of  $\beta$ -crustacyanin are found on irreversible dissociation of  $\alpha$ -crustacyanin at low ionic strength.<sup>83</sup> Whether the carotenoid acts as a direct cross-link between apoproteins, as suggested for the binding of FAD in glucose oxidase<sup>85</sup> and NAD in lactate dehydrogenase,<sup>86</sup> whether it changes the configurations of the apoproteins into those suitable for dimerisation, as for retinol in the binding of retinol-binding protein to prealbumin,<sup>87</sup> or whether it acts as a hydrophobic—Van der Waals bridge, is a matter of conjecture and will be discussed later. Apoproteins of pigments prepared from stored material can no longer give the  $\alpha$ -form of the proteins on recombination with astaxanthin (e.g. for crustacyanin and *A. antennatus* carotenoprotein, Table 5). Prolonged storage similarly inhibits regeneration of the visual pigment, rhodopsin.<sup>88</sup>

The physical properties of crustacyanin and its derivatives have recently been reviewed.<sup>89</sup>

### Compositions

The amino acid compositions of thirteen blue/purple carotenoproteins from diverse tissues of copepod and decapod Crustacea and from two siphonophore species are known. These are listed in a histogram (Fig. 2) in order, from left to right, of decreasing wavelength of absorption maxima. There is no apparent relationship between the content of any particular amino acid or group of amino acids and the position of the absorption maximum of the pigments, except possibly for isoleucine which tends to lower values with decreasing wavelength maximum.

As the size of the carotenoid (*ca.*  $28 \times 5 \text{ \AA}$ ) is comparable with the diameter of the apoprotein (*ca.*  $40 \text{ \AA}$  or less),<sup>71-73</sup> a large part of the protein may be expected to be involved in its binding. If the binding sites are of an unusual nature, these could influence the overall composition. An indication that this may be the case is found when  $S\Delta\phi$  values for pairs of the carotenoproteins are inspected (Table 3). The proteins, from different species and tissue environments, are evidently related in composition. In Fig. 2 the amino acid composition of the "average protein"<sup>93</sup> is given, with the standard error of occurrence of each amino acid as a criterion for comparison. The carotenoproteins are characterised by high contents of acidic (mainly amidated) amino acids, particularly of

aspartic acid. The consistently low contents of leucine and high content of serine, threonine and proline may have some relevance, discussed later, to the binding of carotenoid. The carotenoproteins have a low content of amino acids (glycine and alanine), reflected in the low (anhydrous) mean residue weights. The hydrophobicity values are not particularly high, while the values of the polarity index are similar to those for the soluble proteins and lipoproteins<sup>94</sup> (Table 4).

The copepod and siphonophore carotenoproteins contrast with those of decapods in having fewer aromatic amino acids. In some species tyrosine is either completely absent or present in uncommonly small amounts (Fig. 2). The species investigated inhabit the top few centimetres of the sea surface in tropical waters and are thus exposed to the sun's u.v. radiation. The sensitivity of rhodopsin to bleaching by u.v. light<sup>90</sup> emphasises the possible (cf. 130) advantage to a stable carotenoprotein of a low content of aromatic amino acids. The carotenoprotein of *Veella* also has a high content of cysteine, partly in sulphhydryl form.<sup>31</sup>

### Specificity of the combinations

Examination of the specificity of carotenoid attachment to apocrustacyanin has shown that to effect changes in quaternary structure, the carotenoid must have 4- and 4'-keto groups.<sup>82,83,91</sup> The ability of a range of synthetic and natural carotenoids to bind specifically to apoproteins derived from carotenoproteins of different species has confirmed the earlier conclusion for crustacyanin<sup>91</sup> and extended the studies (Table 5).

(a) *Crustacyanin*. The presence of keto groups in both the 4- and 4'-position of the  $\beta$ -ionone rings (or equivalent positions in the cyclopentene analogues) of the carotenoid is necessary to bring about changes in the structure of the apoprotein, with the formation of  $\beta$ -crustacyanin.

Carotenoids with a 4-keto group in only one  $\beta$ -ionone or cyclopentene ring ( $\beta,\beta$ -carotene-3,4-dione; 3,3',4'-trihydroxy- $\beta,\beta$ -caroten-4-one (idoxanthin); 4'-hydroxy- $\beta,\beta$ -caroten-4-one; 2-nor  $\beta$ -carotene,  $\beta$ -carotene-3,4-dione) show a low affinity for the apoprotein, and apparently fail to bind with sufficient specificity to bring about the configurational changes in the protein necessary for dimerisation.  $\beta,\beta$ -caroten-4-one (echinenone) does not bind to the protein in demonstrable amounts.

Whether the polyenes have planar cyclopentene<sup>92</sup> rings, twisted (relative to the plane of the polyene)  $\beta$ -ionone<sup>72</sup> rings, or a mixture of the two, evidently has no influence on the ability to confer a quaternary structure upon apocrustacyanin. Similar observations have been made with opsin.<sup>98</sup> The absorption bands of the cyclopentene carotenoid-proteins lie at longer wavelengths than those of  $\beta$ -ionone derivatives. Carotenoids with both cyclopentene and  $\beta$ -ionone rings give pigments with two absorption bands, one of which is bathochromically shifted in relation to the spectrum of free carotenoid.

Recombinations obtained with canthaxanthin, astacene and astaxanthin-3,3'-dimethylether show that enolisation of the 4- and 4'-positions is unnecessary and that the 3- and 3'-hydroxyl groups are dispensable for satisfactory recombination. Steric factors or methodology may be responsible for the lack of reconstitution found with astaxanthin diacetate and dipalmitate. Crustaxanthin, in which the shape of both  $\beta$ -ionone rings is altered<sup>79</sup> as compared with astaxanthin, has negligible affinity for the protein; idoxanthin, where the shape of one ring is altered, gives exclusively an apo-sized recombination product, (cf. also 79).

Table 3. Estimation of relatedness ( $S\Delta\phi$  values) of carotenoproteins. Possibly related proteins are in *italic*

Species	Tissue	Decapod carotenoproteins						Siphonophore carotenoproteins	
		<i>A. antennatus</i> Carapace	<i>A. antennatus</i> Stomach	Mandibles	<i>S. arctus</i> Carapace	<i>H. gammarus</i> Carapace	<i>C. erythropus</i> Exoskeleton		<i>E. spinifrons</i> Carapace
<i>A. antennatus</i>	Carapace	0	28	35	38	43	27	64	55
	Stomach	28	0	36	29	48	38	89	78
	Mandibles	35	36	0	60	84	44	76	116
<i>S. arctus</i>	Carapace	38	29	60	0	13	24	89	42
	Carapace	43	48	84	13	0	14	88	17
<i>C. erythropus</i>	Exoskeleton	27	38	44	24	14	0	131	38
	Carapace	64	89	76	89	88	131	0	91
<i>E. spinifrons</i>	Carapace	55	78	116	42	17	38	91	0
	Hypodermis	87	82	93	69	76	94	70	108
<i>Labidocera</i> sp.	Hypodermis	56	89	75	105	116	121	59	149
<i>A. patersoni</i>	Hypodermis	56	88	79	90	96	101	61	128
<i>Verella</i> sp.	Mantle	38	77	63	111	123	130	80	143
<i>Porpita</i> sp.	Mantle and tentacles	70	66	51	105	119	142	66	157

Species	Tissues	Copepod carotenoproteins			Siphonophore carotenoproteins	
		<i>L. acutifrons</i> Hypodermis	<i>Labidocera</i> sp. Hypodermis	<i>A. patersoni</i> Hypodermis	<i>Verella</i> sp. Mantle	<i>Porpita</i> sp. Mantle and tentacle
<i>A. antennatus</i>	Carapace	87	56	56	38	70
	Stomach	82	89	88	77	66
	Mandibles	93	75	79	63	51
<i>S. arctus</i>	Carapace	69	105	90	111	105
	Carapace	76	116	96	123	119
<i>C. erythropus</i>	Exoskeleton	94	121	101	142	130
	Carapace	70	59	61	66	80
<i>E. spinifrons</i>	Carapace	108	149	128	143	157
	Hypodermis	0	80	57	91	89
<i>Labidocera</i> sp.	Hypodermis	80	0	19	42	54
<i>A. patersoni</i>	Hypodermis	57	19	0	49	51
<i>Verella</i> sp.	Mantle	91	42	49	0	9
<i>Porpita</i> sp.	Mantle and tentacle	89	54	51	9	0

Table 4. Comparison of compositions (mol %) of carotenoproteins and different groups of soluble protein, in terms of amino acid sets. Average hydrophobicity and mean (anhydrous) residue weight of carotenoproteins

Carotenoprotein Species	Tissue	Amino acid sets			Small§ (mol %)	charged¶ (mol %)	H $\phi$ avge <sup>  </sup>	M.R.W.††
		Apolar (A)† (mol %)	Polarity index (P)‡ (mol %)	P/A ratio				
<i>A. antennatus</i>	Carapace	19.9	47.6	2.4	20.0	30.7	890	106
	Stomach	16.6	56.0	3.4	15.1	35.3	850	108
	Mandibles	18.5	54.7	3.0	18.3	31.8	790	106
<i>S. arctus</i>	Carapace	20.4	50.8	2.5	14.6	34.6	970	109
<i>C. erythropus</i>	Exoskeleton	19.3	48.5	2.5	15.6	29.7	990	107
<i>E. spinifrons</i>	Carapace	22.1	50.6	2.3	20.9	35.9	780	105
<i>C. clarkii</i>	Carapace	25.8	45.5	1.8	15.8	29.8	1040	106
<i>H. gammarus</i>	Carapace	22.5	46.9	2.1	14.8	31.0	1010	109
<i>L. acutifrons</i>	Hypodermis	25.6	50.1	2.0	16.4	35.0	910	106
<i>Labidocera</i> sp.	Hypodermis	18.8	47.0	2.5	24.4	31.2	810	100
<i>A. patersoni</i>	Hypodermis	20.0	47.8	2.4	22.7	32.2	830	102
<i>Veleva</i> sp.	Mantle	18.5	49.9	2.7	20.4	31.4	780	102
<i>Porpita</i> sp.	Mantle and tentacles	20.7	53.7	2.6	18.8	33.6	770	103
Mean and S.D. for carotenoproteins		20.7 ± 2.7	49.9 ± 3.0	2.5 ± 0.4	18.3 ± 3.3	32.5 ± 3.3		
Soluble lipoproteins <sup>95</sup>		27.3 ± 2.5	50.0 ± 2.4	1.8 ± 0.2	12.8 ± 1.4	33.8 ± 4.0		
Soluble proteins <sup>95</sup>		23.9 ± 3.6	49.1 ± 3.1	2.1 ± 0.5	14.3 ± 4.2	33.3 ± 3.1		
Soluble oligomeric proteins <sup>95</sup>		27.5 ± 2.8	45.1 ± 3.3	1.7 ± 0.6	16.6 ± 3.9	31.0 ± 3.3		

†Apolar residues:<sup>95</sup> Val, Ileu, Leu, Phe, Met.

‡Polarity index:<sup>94</sup> Sum of mol% values of Asp, Thr, Ser, Glu, Lys, His, Arg.

§Small amino acids:<sup>95</sup> Ala, Gly.

¶Charged amino acids:<sup>95</sup> Asp, Glu, Arg, Lys.

<sup>||</sup>Average hydrophobicity, calcs/residues, calculated from hydrophobicity values given in Ref. 96 and corrected values Ref. 97.

††Mean (anhydrous) residue weight.

Acetylenic carotenoids, in which the distance between 4- and 4'-keto groups in the  $\beta$ -ionone or cyclopentene rings is shortened,<sup>85</sup> are capable, with the exception of dehydrocanthaxanthin, of associating the apoprotein units. Mono- and di-acetylenic (7,8 and/or 7',8' positions) carotenoids, pectenoxanthin and pectenolone, occur naturally in the carotenoid-lipoglycoprotein of *Pecten maximus*.<sup>16</sup> Acetylenic (7',8 and/or 7',8' positions) derivatives of astaxanthin are found in the blue carotenoid-protein complex of unknown nature, isolated from the skin of *Asterias rubens*.<sup>99</sup>

The shortened carotenoid 15,15'-dehydro-8'-apo- $\beta$ -carotene-8'al-3, 4-dione ( $\beta$ -apocarotenal) binds to apocrustacyanin without bringing about association of the subunits. Retinals with shortened chains do not form visual pigments<sup>100</sup> or bind to retinol-binding protein of human plasma.<sup>101</sup>

The specificity of the carotenoid-protein association in another decapod carotenoprotein, that of the stomach wall of *A. antennatus*, resembles that in crustacyanin. Carotenoids that have a 4-keto group in one of the  $\beta$ -ionone rings can in some cases (e.g. idoxanthin), however, bring about dimerisation of the apoprotein units. The second  $\beta$ -ionone ring may evidently be accommodated at its binding site even without a 4-keto substituent, provided that there is strong attachment of the first ring. It may be supposed that firm binding of just one  $\beta$ -ionone ring may be sufficient to alter the tertiary, and hence quaternary, structure of the apoprotein. These results imply that the carotenoid does not act as a direct cross-link between the apoproteins. Preliminary recombination studies for the hypodermal pigment of the copepod *A. patersoni* suggests similar specificity.

(b) *Veleva* mantle carotenoprotein. Conflicting reports on the position of the absorption maxima of *Veleva* and

*Porpita* carotenoproteins were attributed previously to a dependence of the carotenoid-protein association on ionic strength and temperature.<sup>102</sup> The presence of specific halide-ion binding sites, which influence the carotenoid-protein interaction, may now be considered as the explanation for the observed variability of the absorption spectra. These sites have been investigated in some detail for the *Veleva* carotenoprotein and are the subject of a forthcoming paper.<sup>31</sup> Briefly, the native carotenoprotein, with absorption maximum at 630 nm, alters on storage in the absence of halide ions to give proteins differing in size, with absorption maxima at 620 nm and 600 nm (in the presence of halide ions). The latter protein, of molecular weight ca. 380,000, was used for the recombination studies. On removal of halide ions this protein, reversibly, both dissociates to purple  $\beta$ -sized units ( $\lambda_{\max}$  570 nm) and alters in configuration to give a derivative with a hypsochromically shifted absorption spectrum ( $\lambda_{\max}$  583 nm) but of unaltered molecular size. The ion-binding sites, three per apoprotein unit, have high (temperature dependent) association constants for the halide ion and probably consist of clusters of basic (arginine and lysine) residues. The spectra for the reconstituted pigments (Table 5) are given in the presence and absence of saturating concentrations of chloride ions.

The allowable alterations in the structure of the carotenoid for pigment formation are similar for both the siphonophore pigment and crustacyanin. The carotenoid derivatives have different effects on the quaternary structures of the two apoproteins. Carotenoids with a 4-keto group in one ring ( $\beta$ -apocarotenal; 4'-hydroxy- $\beta$ , $\beta$ -caroten-4-one; idoxanthin;  $\beta$ , $\beta$ -carotene-3, 4-dione), but not echinenone, bring about dimerisation of the *Veleva* apoprotein. The *Veleva* protein resembles, with exceptions (e.g. 4'-hydroxy- $\beta$ , $\beta$ -caroten-4-one;  $\beta$ , $\beta$ -carotene-



3,4-dione), the *A. antennatus* pigment in this respect.  $\beta$ -Forms of the protein may be obtained when 4-keto groups are present on both rings (e.g. canthaxanthin and astaxanthin diacetate, but not astacene diacetate). Further polymerisation of the protein occurs if the carotenoid has additional hydrophilic substituents (e.g. 2-nor  $\beta$ -carotene,  $\beta$ -carotene-3,4,4'-trione; astacene; violoerythrin), even for acetylenic derivatives (e.g. dehydroastacene and dehydrovioleerythrin).

The *Veleva* apoprotein has two carotenoid-, and so four  $\beta$ -ionone-, binding sites.<sup>31</sup> One can envisage that the 4-monoketo carotenoids are bound so that the 4-keto- $\beta$ -ionone groups occupy two sites on one face of the protein and provide the correct configuration along that face (only) for interaction with a similar complementary face of a second apoprotein unit; polymerisation cannot then proceed further than the  $\beta$ -form of the protein.

(c) *Ovorubin*. The length of the carotenoid, the size and shape of the  $\beta$ -ionone ring and the angle of the ring about the 6-7 bond to the plane of the polyene chain can be altered without inhibiting binding of the carotenoid. Enolisation of the 4- and 4'-keto groups, or of the 3- and 3'-keto groups, is not required for carotenoid binding since  $\beta$ -apocarotenal, astaxanthin diacetate, violoerythrin and dehydrovioleerythrin can give pigments with apovorubin. The binding nonetheless is fairly selective since canthaxanthin and 2-nor  $\beta$ -carotene,  $\beta$ -carotene-3,4-dione have no affinity for the binding site, in contrast to 4'-hydroxy- $\beta$ , $\beta$ -caroten-4-one and  $\beta$ , $\beta$ -carotene-3,4-dione.

The nature of the binding site, which alters the relative importance of excitations to different vibrational levels in the associated carotenoid giving rise to the fine structure seen in the spectrum, is unknown. Ovorubin can be dissociated with sodium dodecylsulphate and separated into two glycoprotein fractions with molecular sizes of about  $2.2 \times 10^4$  and  $3.5 \times 10^4$ , respectively. The larger fraction recombines with astaxanthin to give the characteristic absorption spectrum of the original pigment.<sup>103</sup>

#### Mode of binding

It has been emphasised that in the spectrum of crustacyanin the extinction coefficient, half-band width and oscillator strength of astaxanthin are not significantly altered compared with that of the carotenoid dissolved in hexane.<sup>79</sup> The same is true for the carotenoproteins listed in Table 6 and for some other carotenoproteins (cf. half-band width of *Pontella fera*<sup>14</sup> and *Cambarus clarkii*<sup>73</sup> carotenoproteins and some decapod stomach carotenoproteins<sup>15</sup>). The invariability of the shape of the absorption spectra, in marked contrast to the alterations in the spectra of the bound carotenoid in carotenoid-lipoglycoproteins, implies that a common binding mechanism exists for most carotenoproteins. It has been stressed that it is the over-all excitation energy of astaxanthin that is altered in crustacyanin, and that there is no large change in the nature of the electronic transition and relative importance of excitation to different vibrational levels. Polarisation of the carotenoid, charge transfer interaction and medium effects thus cannot explain the bathochromic shift in the absorption spectrum of the bound astaxanthin.<sup>79</sup>

In crustacyanin, and other carotenoproteins, astaxanthin is presumably bound non-covalently, since it is easily removed by organic solvents. No evidence has been obtained for labile Schiff's base formation between the keto groups of astaxanthin and amino groups of crustacyanin, or of linkages involving sulphhydryl groups.<sup>79,81</sup> A sugges-

tion<sup>72</sup> that astaxanthin may be bound by unstable anhydride bonds to the carboxyl groups of aspartic acid has been disproved experimentally. Amidation of the carboxyl groups of apocrustacyanin does not eliminate the affinity of the protein for its prosthetic group.<sup>51</sup> Non-covalent electrostatic attachment, originally suggested by Kuhn and Sørensen to account for the colour of ooverdin, in which the bis-dianion of astaxanthin is bound to basic groups of the protein, is unlikely to be the mode of binding in crustacyanin both from spectroscopic evidence<sup>79</sup> and from the ability of astacene, canthaxanthin and violoerythrin to give coloured products with the apoprotein. Recently the astaxanthin removed from crustacyanin (and ooverdin) has been shown to be optically active in the 3- and 3'-position (3S, 3'S)<sup>104</sup> which is difficult to explain if it is bound as the tetra-anion. The low content of hydrophobic amino acids and the not unusual average hydrophobicity values for the carotenoproteins, would seem to make hydrophobic bonding, of the type suggested for retinol in retinol-binding protein,<sup>105-7</sup> unlikely. In addition, studies in the effects of solvents on the spectrum of astaxanthin showing the insensitivity of the absorption band to the nature of the solvent,<sup>82</sup> make it improbable that hydrophobic bonding alone could account for the large bathochromic shifts in the absorption bands of the pigments. The failure to show induced fluorescence of the carotenoid in crustacyanin upon excitation at 280 nm,<sup>79</sup> in contrast to the results for retinol-binding protein,<sup>108</sup> would indicate that aromatic amino acids are probably absent from the vicinity of the polyene. Exciton interaction between carotenoid molecules, aligned head to tail, on adjacent apoprotein units is an improbable reason for the spectral shifts of the bound carotenoid in carotenoproteins as it would alter the shape of the spectrum.<sup>79</sup> It could, however, explain what appears to be a splitting in the absorption band in the spectrum of *Labidocera* sp. carotenoprotein (Table 6).

A model, based on chemical analogies, which satisfactorily accounts for the absorption spectra of blue and purple carotenoproteins, has been put forward by Buchwald and Jencks.<sup>79</sup> It is proposed that the spectral characteristics of the crustacyanin pigments (and rhodopsin)<sup>79,109</sup> are due to distortion of the polyene about the double bonds with a consequent decrease in double-bond order. For the mechanism to be valid, the methyl groups of the polyene chain must be immobilised and the  $\beta$ -ionone rings bound to the protein so as to be twisted out of the plane of the conjugated polyene, thereby localising the strain in the double, rather than single, bonds. The shape of the absorption band in the visible region should then be unaltered but bathochromically shifted by an amount depending on the degree of twist. The theory is consistent with the lack of demonstrable intermediates during the denaturation of crustacyanin,<sup>79</sup> and for the more strongly bathochromically shifted absorption bands of analogous pigments formed from flat cyclopentene carotenoid derivatives (e.g. violoerythrin and dehydrovioleerythrin, Table 5). The more complex spectra of the mixed cyclopentene,  $\beta$ -ionone ring carotenoproteins may mean that in these combinations additional polarisation effects are superimposed on the twisting. In the case of rhodopsin, the binding is known to involve the protonated Schiff's base of the polyene; medium and counter-ion effects, as well as twisting of the chromophore, have been suggested as causes for the bathochromic shift in the absorption spectrum.<sup>110</sup> The ability of 9,13-desmethylretinal and other demethylated














4'-Hydroxy- $\beta,\beta$ -caroten-4-one		9	5% [510-515]	95% 492( $\pm$ Cl $^-$ )	13	565	Nil	Trace	[460]	19	495
$\beta,\beta$ -carotene-3,4-dione (euglenanone)		11	75% 517	25% 495	11	550	Nil	Trace	[450]	6	465-470
4'-Hydroxy- $\beta,\beta$ -carotene-3,4-dione		Nil									
3,3',4'-Trihydroxy- $\beta,\beta$ -caroten-4-one (floxanthin)		16	90% 535( $\pm$ Cl $^-$ )	10% 515( $\pm$ Cl $^-$ ) <sup>†</sup>	18	563	25	Trace	[475]	36	450
$\beta,\beta$ -Carotene-4,4'-dione (canthaxanthin)		2	15% 560-570	85% 510	12	575	8	Trace	[580,485]	Nil	
15,15'-Dehydrocanthaxanthin		3		485	6	545	Trace	Trace	[470-480]	Nil	
3,3'-Dihydroxy- $\beta,\beta$ -carotene-4,4'-dione (astaxanthin)			90% 600(+Cl $^-$ ) 580(-Cl $^-$ )	2% 545							485 510 545
Astaxanthin-3,3'-diacetate		6 $\ddagger$	10% 560-570	90% 510	Nil					10	(490) 515 560

Table 5(Contd)

Carotenoid	Veilella Carotenoprotein <sup>(3)</sup>		Crustacyanin		A. Antennatus (stomach) carotenoprotein <sup>(4)</sup>		A. Patersoni carotenoprotein		Ovonubin <sup>(5)</sup>	
	Excluded from Sephadex G75	Absorption maxima (nm)	"β-Size"	"Apo-size"	Absorption maxima <sup>(2)</sup> (nm)	"β-Size"	"Apo-size"	Absorption maxima (nm)	Absorption maxima (nm)	Absorption maxima (nm)
Astaxanthin-3,3'-dimethylether	Nil				565	Nil†		Trace	Nil	470
Astaxanthin-3,3'-diacrylate§	Trace					Nil†			5	
Astaxanthin-3'-acrylate§										
β,β-carotene-3,3',4,4'-tetraone (astacene)										
										
Astacene-3,3'-diacetate	Nil†								Nil	460
15,15'-Dehydroastacene	14	80% 552 (+Cl <sup>-</sup> ) 537 (-Cl <sup>-</sup> )	10% 490	10% 537 (+Cl <sup>-</sup> ) 515 (-Cl <sup>-</sup> )	555	7	528	Trace	10	490 520
3,3',4,4'-Tetrahydroxy-β,β-carotene (crustaxanthin)	3		(450) 472 (505)						6	480 510
										
2-Nor β-carotene,β-carotene-3,4'-dione	6	95% 625,405 (±Cl <sup>-</sup> )								
										
2-Nor β-carotene,β-carotene-3,4,4'-trione	10	98% 725,(480-490) (+Cl <sup>-</sup> ) 695,(480-490) (-Cl <sup>-</sup> )								
										

Undet†  
(>10%)  
Nil  
Nil†

Nil†

70

575

30

Nil†

1%

555

7

528

Trace

10

480  
490  
520

65% (±Cl<sup>-</sup>)  
505 (±Cl<sup>-</sup>)

25%  
535 (+Cl<sup>-</sup>)  
520 (-Cl<sup>-</sup>)

10%  
537 (+Cl<sup>-</sup>)  
515 (-Cl<sup>-</sup>)

635  
(415)

635  
590  
510

650§§  
590  
510

695,510

Trace

15

695,510

9

553  
(530)

†

†

†

†

†

†

†

†

†

†

†

†

15,15'-Dehydro-2-nor-β-carotene-β-carotene-3,4,4'-trione	32	95% 660, 455 +Cl <sup>-</sup> 630, 440 -Cl <sup>-</sup>	5% 640, 440 +Cl <sup>-</sup> 627, 430 -Cl <sup>+</sup>	21	635, 485	30	630, 460	‡	[660]	10	(485) 515 560	
2,2'-Dinor-β,β-carotene-3,3',4,4'-tetraone (violerythrin)	14	15% 670 (+Cl <sup>-</sup> ) 650 (-Cl <sup>-</sup> )	75% 645 (+Cl <sup>-</sup> ) 635 (-Cl <sup>-</sup> )	10% [520]	8	636	5	[650]	‡	[670]	4	(480) 510 560
15,15'-Dehydroviolerythrin	41	75% 670 (+Cl <sup>-</sup> ) 645 (-Cl <sup>-</sup> )	23% 650 (+Cl <sup>-</sup> ) 637 (-Cl <sup>-</sup> )	2%	75	670	30	655, (440)	‡	[680]	27	(485) 520 550
3,3'-Dihydroxy-2,2'-dinor β,β-carotene-4,4'-dione-3,3'-diacrylate (actinoerythrin)	Trace			Nil†							Nil	
15,15'-Dehydro-8'-apo-β-carotene-8'-al-3,4'-dione	51		95% 508 (±Cl <sup>-</sup> )	3	504	7	Trace	450-480	Trace	[500]	25	465
Carotenoid (unidentified) isolated from the carotenoid-protein complex of the muscle tissue of <i>Salpa cylindrica</i>	2		470-490	Nil		Nil						

1. % Recombinations are derived from the extinction of the complexes at the main absorption maximum, assuming identical extinction coefficients and taking the recombination obtained with astaxanthin as 100%. Carotenoid-protein complexes are separated in gel filtration using Sephadex G75 into protein excluded from the gel, protein having a similar elution volume to the β-form of the native pigment, and that showing a similar size to the apoprotein.

2. Spectral data are for the proteins dissolved in 0.1 M phosphate buffer, pH 7. Inflections are given in brackets. Values in square brackets are approximate.

3. *Veillela* carotenoprotein (λ max 600 nm) was used in the studies. Absorption maxima of the complexes dissolved in 0.1 M phosphate buffer, pH 7, are recorded at 5°C in the presence and, where stated, in the absence of 0.2 M KCl.

4. The sizes of reconstituted pigments were not investigated.

5. Absorption maxima recorded for those carotenoids associating feebly with apoprotein are approximate; the spectra are partially obscured by traces of strongly bound helicorubin present in the freeze-dried overubin used in the studies.

†Less than 10% recombination (based on 100% recombination with astaxanthin).

‡Greater than 10% recombination.

§Mixture of long-chain fatty acids.

||Results taken from Ref. 83.

¶Isolated pigments contained astaxene.

¶¶Carotenoid-protein complexes observed initially were not apparent following gel filtration.

††"Apo-size" material contains astaxanthin and astaxanthin diacetate.

§§Absorption maxima prior to gel filtration.

Table 6. Characteristics of the absorption spectra of carotenoproteins

Species	Tissue	$\lambda_{\max}$ † (nm)	$V_{\frac{1}{2}}$ (cm <sup>-1</sup> )‡ $\times 10^{-2}$	$E_{\max}$ (cm <sup>-1</sup> )§ $\times 10^{-5}$	f¶
Aristeus antennatus <sup>71</sup>	Carapace	593	43.0	1.23	2.46
	Mandibles	595	44.0		2.32 <sup>  </sup>
	Stomach	588	43.0	1.24	2.56
Palinurus vulgaris <sup>71</sup>	Carapace	560	45.0		2.67 <sup>  </sup>
Scyllarus arctus <sup>71</sup>	Carapace	616	46.0	1.20	2.70
Clibanarius erythropus <sup>71</sup>	Exoskeleton	620	41.0	1.26	2.34
Galathea strigosa <sup>71</sup>	Carapace	589	46.0		2.64 <sup>  </sup>
Eriphia spinifrons <sup>71</sup>	Carapace	536	42.0		2.50 <sup>  </sup>
Homarus gammarus <sup>79</sup> (L.)	Exoskeleton	632	42.0	1.25	2.60
Labidocera acutifrons <sup>72</sup>	Hypodermis	640	42.0	1.19	2.52
Labidocera sp. <sup>31</sup>	Hypodermis	630, 660	43.0	1.26	2.54
Anomalocera patersoni <sup>31</sup>	Hypodermis	650	44.0	1.26	2.46
Verella sp.††	Mantle	620	40.0	1.28	2.63
Verella sp.	Mantle	600	39.0	1.27	2.38
Porpita sp. <sup>31</sup> ††	Mantle & tentacles	650	43.4	1.28	2.60
Astaxanthin					
In pyridine <sup>79</sup>		492	42.0	1.12	2.35
In hexane <sup>79</sup>		472	42.0	1.24	2.60

†Position of main absorption maximum at room temperature for the pigment dissolved in 0.2 M phosphate buffer (KH<sub>2</sub>PO<sub>4</sub>-Na<sub>2</sub>HPO<sub>4</sub>), pH 7.

‡Half-band width.

§Molar extinction coefficient, based on astaxanthin content.

¶Oscillator strength.

<sup>||</sup>Assuming a value of  $1.23 \times 10^5$  for  $E_{\max}$ .

††Carotenoprotein dissolved in 1 M KCl-0.05 M phosphate buffer, pH 7.

derivatives of retinal to produce visual pigments in which the absorption maxima of the polyenes are still strongly displaced to longer wavelength,<sup>98,100</sup> is difficult to reconcile with the twisting mechanism.

The strong optical activity of the bound astaxanthin in  $\alpha$ - and  $\beta$ -crustacyanin,<sup>79</sup> and of retinal in rhodopsin and retinochrome,<sup>112</sup> has similarly been attributed to extrinsic stereospecific twisting of the polyenes which introduces asymmetry into the chromophores. The magnitude of the induced optical activity of retinal in rhodopsin can equally well be explained by a Kirkwood-type coupled oscillator mechanism, involving the interaction of the polyene transition with the transition of a neighbouring (< 15 Å) aromatic side chain.<sup>98,110,113</sup> The ORD and circular dichroism (CD) spectra of the crustacyanins in the visible region exhibit evidence of degenerate exciton interaction. The CD spectrum of  $\beta$ -crustacyanin has negative and positive Cotton effects above and below the position of the absorption maximum, respectively.<sup>79</sup> The splitting of the excitation may be the result of intermolecular interaction of the transition dipoles of carotenoids at an angle to one another and attached to closely situated apoprotein units.<sup>79,82</sup> An alternative explanation, previously suggested for the exciton splitting of the CD spectrum of *N*-all-*trans*-retinylidene-poly-L-lysine,<sup>114</sup> could be mutual induction of optical activity in the chromophores due to an asymmetrical arrangement of the (adjacent) binding sites. The *Verella* pigment, the only other carotenoprotein of which the optical activity has been investigated, resembles crustacyanin in showing a large maximal molar ellipticity value for the bound carotenoid. Splitting of the excitation, inverted in shape compared with that in crustacyanin, is

seen only with the form of the protein with absorption maximum at 620 nm and only in the presence of chloride ions (Fig. 3);<sup>31</sup> there appears to be no major change in the protein conformation in this process, as revealed in the u.v. CD spectra.




Fig. 3. Circular dichroism spectrum of *Verella* carotenoprotein ( $\lambda_{\max}$  620 nm) dissolved in 0.2 M phosphate (KH<sub>2</sub>PO<sub>4</sub>-Na<sub>2</sub>HPO<sub>4</sub>) buffer, pH 7, in the absence (A) and presence of (B) of 0.5 M KCl.

The absorption spectra of  $\alpha$ - and  $\beta$ -crustacyanin, broad and featureless at room temperature, sharpen somewhat at 77°K to reveal incipient vibrational structure. The spectrum of astaxanthin, the chromophore, sharpens markedly at 77°K to reveal a typical vibrational progression of  $1270 \pm 60 \text{ cm}^{-1}$ .<sup>115</sup> Although the circular dichroism spectra of  $\alpha$ - and  $\beta$ -crustacyanin show double-peaked degenerate exciton interaction under the long-wavelength long-axis polarised band, no splitting is visible under the short-axis polarised *cis* peak at *ca.*  $2700 \text{ cm}^{-1}$ .<sup>115</sup> In principle geometrical information about the relative disposition of a pair of carotenoids can be extracted from the spectrum of  $\beta$ -crustacyanin. In practice this is difficult to achieve unambiguously. However, the lack of degenerate exciton splitting under the *cis* peaks is evidence that the directions of these two short axis transition dipole moments on interacting carotenoids must be parallel. Thus the planes containing the long axis of each carotenoid must also be parallel. Further analysis of the degenerate exciton interaction under the long wavelength band gave an angle of *ca.* 77° between the two long axes of the carotenoids. The distance between their centres was not obtained with certainty but must lie in the range 4–10°Å in order to give effects of the magnitude observed. In the aggregated  $\alpha$ -derivative a bond angle of 72° between the long axes was deduced.<sup>115</sup> At these distances the London dispersion—Van der Waals forces between two oriented carotenoids<sup>116</sup> may be of sufficient magnitude to make an appreciable contribution to the dimerisation of the apoprotein units.

#### Nature of the carotenoid-binding sites

One can only speculate on the nature of the carotenoid-binding sites in carotenoproteins. The carotenoid is believed to be buried in the association, removed from water,<sup>78</sup> but with the 4- and 4'-keto groups near the protein surface.<sup>79</sup> The proposed strain mechanism for the binding requires close fit between the polypeptide and methyl groups of the polyene chain and firm anchoring of the  $\beta$ -ionone rings.<sup>79</sup> The high content of small amino acids, particularly in proteins of low minimum molecular weight (based on astaxanthin contents), may facilitate an intimate fit between polyene and polypeptide backbone. It is probable that the structures of carotenoproteins are composed mainly of random coil and  $\beta$ -conformations. The low content of leucine, the main stabilising influence in the inner sections of  $\alpha$ -helices,<sup>117</sup> and the high content of helix-breaking<sup>118,9</sup> amino acids (proline, serine, glycine and asparagine) indicate that the proteins must have low helical contents. Confirmation of this, for crustacyanin and *Veillella* carotenoprotein, has come from u.v. CD measurements. The helical content of crustacyanin has been estimated to be 6%.<sup>79</sup> The *Veillella* carotenoprotein may have a configuration composed almost entirely of  $\beta$ -structure.<sup>31</sup> Amino acids which occur in high frequency in  $\beta$ -structures (isoleucine, valine, threonine and glutamine)<sup>117,121</sup> are not infrequent in carotenoproteins. A large number of  $\beta$ -bends<sup>120</sup> in the structures, assisting the formation of anti-parallel  $\beta$ -pleated sheets may result from the high contents of aspartic acid, serine, threonine and proline. The repeat distance between similarly oriented groups (6.68 Å)<sup>122</sup> in  $\beta$ -pleated sheet structures is of similar magnitude to the distance between methyl groups in the polyene chain.<sup>123</sup> It is suggested that immobilisation of these methyl groups occurs through binding to lengths of  $\beta$ -pleated sheet in which small substituents (e.g. alanine) favour strong hydrophobic

interaction, as exists between polyene chains in canthaxanthin crystals.<sup>123</sup> The  $\beta$ -ionone ring, twisted out of the plane of the polyene chain about the 6–7 single bond, may interact hydrophobically through a methyl group in the 1-position with a flexible residue (e.g. isoleucine) of the  $\beta$ -pleated sheet. It is also proposed, in view of the importance of 4-keto groups for combination with apoprotein, that the  $\beta$ -ionone rings are bound through these groups by hydrogen-bonding to imide groups of peptide bonds on peripheral chains of  $\beta$ -pleated sheet structures (Fig. 4). The hydrogen bonding between peripheral and adjacent chains of the  $\beta$ -structure would ensure, through electron withdrawal, a stronger hydrogen-bonding of the carotenoid than would be obtained with the single chain. The 5- and 5'-methyl groups of the prosthetic group may be made secure by bonding to an adjacent hydrophobic residue (e.g. isoleucine). The binding can be additionally stabilised, as molecular models show (preferred hydrogen-bond angles and distances<sup>124</sup> being maintained) by hydrogen-bonding of the 3S- and 3'S-hydroxy groups of the carotenoid to the neighbouring peptide bond carbonyl groups. The relative orientation of the pieces of  $\beta$ -structure binding the carotenoid would then control the twisting of the polyene and consequently the absorption maximum of the pigment.

#### BACTERIAL AND PLANT CAROTENOID-PROTEIN COMPLEXES AND VITAMIN A-PROTEINS

A number of lipoprotein particles or complexes containing chlorophyll and/or cytochromes and non-covalently bound carotenoids, in which there is some selectivity in the nature of the associated carotenoids, have been isolated from membranes of plant chloroplasts.<sup>125–9</sup> The complexity of the particles precludes any meaningful discussion as to the mode of binding of the carotenoid component, which may involve protein, lipid and chlorophyll.<sup>133</sup> The preparation and properties of conjugates of  $\beta$ -carotene with insoluble<sup>131</sup> and soluble<sup>134</sup> chloroplast lamellar(lipo)protein, free from other components, have recently been reviewed. It is still controversial whether these are natural complexes, or artifacts. These and other plant "carotenoproteins"<sup>132,135</sup> may be various types of carotenoid aggregates in stabilised suspension,<sup>82,134,136,137</sup> or may be formed by unspecific association of carotenoid with hydrophobic protein during isolation (cf. 107, 144).




Fig. 4. Proposed mode of binding of astaxanthin in carotenoproteins.

Table 7. Comparison of compositions (mol %) of bacterial and plant carotenoid-protein complexes and vitamin A-proteins, in terms of amino acid sets, with those of membrane lipoproteins and different groups of soluble proteins

Protein	Apolar residues (A)† (mol %)	Polarity index (P)‡ (mol %)	P/A ratio	Small§ (mol %)	Charged¶ (mol %)
Rhodopsin (bovine) <sup>151</sup>	29.3	39.9	1.4	18.0	22.7
Rhodopsin (treated with <sup>147</sup> chymotrypsin)	36.2	37.7	1.0	16.6	23.9
N-retinylidene peptide <sup>148</sup>	40.0	20.0	0.5	30.0	—
Bacteriorhodopsin <sup>152</sup>	34.8	35.5	1.0	21.4	23.1
Spirilloxanthin-glycolipoprotein <sup>139</sup> complex ( <i>Rhodospirillum rubrum</i> )	29.4	41.1	1.4	22.3	28.7
Detergent-soluble carotenoid- lipoprotein complex ( <i>Sarcina</i> <sup>141</sup> <i>flava</i> )	38.2	38.4	1.0	19.4	16.7
Carotenoidglycoside-peptide ( <i>S. flava</i> , fraction 4 (iii)), ether- soluble following saponification step <sup>142</sup>	19.9	48.6	2.4	28.7	33.5
Bacteriuberterglycoside-peptide ( <i>Sarcina morrhuae</i> ), water-soluble following saponification step <sup>143</sup>	28.6	44.7	1.6	24.1	35.3
$\beta$ -carotene-binding lamellar protein (spinach chloroplast), water-soluble <sup>134</sup>	31.0	39.7	1.3	20.1	27.7
$\beta$ -carotene-binding lamellar protein (spinach chloroplast), water-insoluble <sup>133</sup>	31.0	39.5	1.3	20.1	27.7
$\beta$ -carotene-containing cytochrome b <sup>559</sup> (spinach chloroplast) <sup>129</sup>	33.0	39.5	1.2	17.6	18.5
Retinol-binding protein (serum) (average composition: rat, porcine, monkey, human) <sup>150</sup>	25.5	51.1	2.0	13.0	37.6
Soluble lipoproteins <sup>95</sup>	27.3 $\pm$ 2.5	50.0 $\pm$ 2.4	1.8 $\pm$ 0.2	12.8 $\pm$ 1.4	33.8 $\pm$ 4.0
Soluble proteins <sup>95</sup>	23.9 $\pm$ 3.6	49.1 $\pm$ 3.1	2.1 $\pm$ 0.5	14.3 $\pm$ 4.2	33.1 $\pm$ 3.1
Soluble oligomeric proteins <sup>95</sup>	27.5 $\pm$ 2.8	45.1 $\pm$ 3.3	1.7 $\pm$ 0.6	16.6 $\pm$ 3.9	31.0 $\pm$ 3.3
Membranous lipoproteins <sup>95</sup>	29.9 $\pm$ 2.8	42.7 $\pm$ 5.7	1.4 $\pm$ 0.3	16.9 $\pm$ 3.0	28.8 $\pm$ 5.8

† Apolar residues:<sup>95</sup> Val, Ileu, Leu, Phe, Met.

‡ Polarity index:<sup>94</sup> sum of mol % values of Asp, Thr, Ser, Glu, Lys, His, Arg.

§ Small amino acids:<sup>95</sup> Ala, Gly.

¶ Charged amino acids:<sup>95</sup> Asp, Glu, Arg, Lys.

Functional particles containing bacteriochlorophyll and non-covalently bound carotenoids have, likewise, been isolated from the chromatophores of photosynthetic bacteria.<sup>138</sup> The spectra of carotenoid-protein complexes, freed from bacteriochlorophyll derived from *Rhodospirillum rubrum* particles, are hypsochromically shifted from that of the carotenoid in the native or free state<sup>131,138-40</sup> (cf. the spectra of some plant<sup>135</sup> and invertebrate<sup>33-6,82</sup> carotenoid-protein complexes). The low molecular weight and the low carotenoid content of the purified *R. rubrum* spirilloxanthin-glycolipoprotein,<sup>140</sup> make it unlikely that this complex is a carotenoid aggregate.<sup>131</sup> The relationship between the state of the carotenoid in the complex and that *in vivo*, however, is yet to be established. An unspecific combination is not excluded.

Detergent-soluble complexes, in which a carotenoid glucoside is covalently bound to lipoprotein, and ether-soluble and water-soluble carotenoidglycoside-peptides have been isolated from the membranes of non-photosynthetic bacteria and partially characterised.<sup>141-3</sup>

Recent reviews and articles on the retinol-binding protein of serum,<sup>101,145,153</sup> retinochrome,<sup>112,146</sup> bacteriorhodopsin<sup>147</sup> and the spectra of visual pigments<sup>110</sup> are relevant to the present review insofar as they throw light on polyene-protein interactions.

Most of the proteins considered in this section are part of membrane structures and have amino acid composi-

tions characteristic of membrane proteins, often with high contents of small amino acids. The retinol-binding protein has an overall composition similar to those of typical soluble lipoproteins, while the carotenoidglycoside-peptides of *Sarcina flava* and *Sarcina morrhuae* bear some resemblance in the different groups of amino acids to the invertebrate carotenoproteins (Table 7).

*Acknowledgements*—I am deeply indebted to Professor D. F. Cheesman (Bedford College) for invaluable advice and assistance during the preparation of the review and to Dr. B. M. Gilchrist (Bedford College) for kindly criticising the manuscript. I would like to thank Dr. B. M. Gilchrist, P. J. Herring (National Institute of Oceanography, Surrey) and A. J. Thomson (University of East Anglia) for allowing me to mention their unpublished observations. I am grateful to Professor B. C. L. Weedon (Queen Mary College, London) for kindly supplying synthetic carotenoids and to Dr. P. M. Scopes (Westfield College, London) for measuring the CD spectra of the *Veillela* pigments.

*Addendum*—In a recent paper (*J. Exp. Zool.* **188**, 289-296 (1974)) C. W. Lui, B. A. Sage and J. D. O'Connor have demonstrated that ovaries of *Procambarus* sp. are capable, *in vitro*, of forming at least part of the lipovitellin molecule.

#### REFERENCES

- D. F. Cheesman, W. L. Lee and P. F. Zagalsky, *Biol. Rev.* **42**, 132 (1967).
- H. Thommen, *Carotenoids* (editor O. Isler) Chap. 8. Birkhäuser, Basel (1971).



- <sup>3</sup>D. Bellamy, *Comp. Biochem. Physiol.* **17**, 1137 (1965).
- <sup>4</sup>D. G. Cornwell, F. A. Kruger and H. B. Robinson, *J. Lipid Res.* **3**, 65 (1962).
- <sup>5</sup>M. Fingerman, *Physiol. Rev.* **45**, 296 (1965).
- <sup>6</sup>T. W. Goodwin, *The Physiology of Crustacea* (editor T. H. Waterman), Vol. 1, Chap. 3, Academic Press, New York (1960).
- <sup>7</sup>R. Elofsson and E. Hallberg, *J. Ultrastruct. Res.* **44**, 421 (1973).
- <sup>8</sup>D. L. Fox, *Comp. Biochem. Physiol.* **44B**, 953 (1973).
- <sup>9</sup>P. J. Herring, *J. Marine Biol. Assoc. UK* **53**, 539 (1973).
- <sup>10</sup>D. L. Fox and T. S. Hopkins, *Comp. Biochem. Physiol.* **17**, 841 (1966).
- <sup>11</sup>D. L. Fox and T. S. Hopkins, *Comp. Biochem. Physiol.* **19**, 267 (1966).
- <sup>12</sup>D. L. Fox and D. W. Wilkie, *Comp. Biochem. Physiol.* **36**, 49 (1970).
- <sup>13</sup>D. L. Fox, *Comp. Biochem. Physiol.* **43B**, 919 (1972).
- <sup>14</sup>P. J. Herring, *Symp. zool. Soc. Lond.* **19**, 215 (1967).
- <sup>15</sup>P. J. Herring, *J. Marine Biol. Assoc. UK* **52**, 179 (1972).
- <sup>16</sup>P. F. Zagalsky, D. F. Cheesman and H. J. Ceccaldi, *Comp. Biochem. Physiol.* **22**, 851 (1967).
- <sup>17</sup>B. M. Gilchrist and W. L. Lee, *Comp. Biochem. Physiol.* **42B**, 263 (1972).
- <sup>18</sup>D. L. Fox, T. S. Hopkins and D. B. Zilversmit, *Comp. Biochem. Physiol.* **14**, 641 (1965).
- <sup>19</sup>G. C. Chen and J. P. Kane, *Biochemistry* **13**, 3330 (1974).
- <sup>20</sup>P. J. Herring, *Proc. 9th Europ. Marine Biol. Symp.* 299 (1975).
- <sup>21</sup>P. J. Herring, unpublished observations.
- <sup>22</sup>D. L. Fox, V. E. Smith and A. A. Wolfson, *Experientia* **23**, 12 (1967).
- <sup>23</sup>A. R. Holter, *Comp. Biochem. Physiol.* **28**, 675 (1968).
- <sup>24</sup>A. H. Warner, J. G. Puodziukas and F. J. Finamore, *Exp. Cell Res.* **70**, 365 (1972).
- <sup>25</sup>P. F. Zagalsky and B. M. Gilchrist, *Comp. Biochem. Physiol.*, In press.
- <sup>26</sup>D. F. Cheesman and J. P. Prebble, *Comp. Biochem. Physiol.* **17**, 929 (1966).
- <sup>27</sup>W. Fyffe and J. D. O'Connor, *Comp. Biochem. Physiol.* **47B**, 851 (1974).
- <sup>28</sup>H. J. Ceccaldi, D. F. Cheesman and P. F. Zagalsky, *Compt. Rend. Soc. Biol.* **160**, 587 (1966).
- <sup>29</sup>D. L. Fox, V. Elliott Smith and A. A. Wolfson, *Experientia* **23**, 965 (1967).
- <sup>30</sup>D. A. Norden, *Comp. Biochem. Physiol.* **42B**, 569 (1972).
- <sup>31</sup>P. F. Zagalsky and P. J. Herring, Papers in preparation.
- <sup>32</sup>B. M. Gilchrist, Unpublished observations.
- <sup>33</sup>B. M. Gilchrist, *Comp. Biochem. Physiol.* **24**, 123 (1968).
- <sup>34</sup>W. L. Lee, *Comp. Biochem. Physiol.* **18**, 17 (1966).
- <sup>35</sup>W. L. Lee and B. M. Gilchrist, *J. exp. Marine Biol. Ecol.* **10**, 1, (1972).
- <sup>36</sup>W. L. Lee, *Comp. Biochem. Physiol.* **19**, 13 (1966).
- <sup>37</sup>J. Green, *Biol. Rev.* **40**, 580 (1965).
- <sup>38</sup>J. Green, *J. Zool. Lond.* **147**, 174 (1966).
- <sup>39</sup>P. J. Herring, *Comp. Biochem. Physiol.* **24**, 205 (1968).
- <sup>40</sup>P. J. Herring, *Comp. Biochem. Physiol.* **24**, 187 (1968).
- <sup>41</sup>R. A. Wallace, S. L. Walker and P. V. Hauschka, *Biochemistry* **6**, 1582 (1967).
- <sup>42</sup>F. Leuenberger and H. Thommen, *J. Insect Physiol.* **16**, 1855 (1970).
- <sup>43</sup>R. A. Wallace, *Biochim. Biophys. Acta* **74**, 505 (1963).
- <sup>44</sup>J. L. Oncley, F. R. W. Gurd and M. Melin, *J. Am. chem. Soc.* **72**, 458 (1950).
- <sup>45</sup>R. A. Wallace, *Analyt. Biochem.* **11**, 297 (1965).
- <sup>46</sup>H. J. Ceccaldi, Thèse de doctorat ès-sciences naturelles, Université d'Aix-Marseille (1968).
- <sup>47</sup>C. Jacobs and W. P. Jencks, *Biol. Bull.* **129**, 410 (1965).
- <sup>48</sup>E. W. Bergink and R. A. Wallace, *J. Biol. Chem.* **249**, 2897 (1974).
- <sup>49</sup>E. W. Bergink, R. A. Wallace, J. A. Van de Berg, E. S. Bos, M. Gruber and A. B. Geert, *Amer. Zool.* **14**, 1177 (1974).
- <sup>50</sup>R. K. Dejmaj and V. J. Brookes, *J. Biol. Chem.* **247**, 869 (1972).
- <sup>51</sup>P. F. Zagalsky, unpublished observations.
- <sup>52</sup>R. Smith, J. R. Dawson and C. Tanford, *J. Biol. Chem.* **247**, 3376 (1972).
- <sup>53</sup>W. H. Cook and W. G. Martin, *Structural and Functional Aspects of Lipoproteins in Living Systems* (editors E. Tria and A. M. Scanu), Chap. C8. Academic Press, New York (1969).
- <sup>54</sup>A. Scanu, *Structural and Functional Aspects of Lipoproteins in Living Systems* (editors E. Tria and A. M. Scanu), Chap. C3. Academic Press, New York (1969).
- <sup>55</sup>S. Margolis, *Structural and Functional Aspects of Lipoproteins in Living Systems* (editors E. Tria and A. M. Scanu), Chap. C2. Academic Press, New York (1969).
- <sup>56</sup>P. F. Zagalsky, *Comp. Biochem. Physiol.* **41B**, 385 (1972).
- <sup>57</sup>J. J. Marchalonis and J. K. Weltman, *Comp. Biochem. Physiol.* **38B**, 609 (1971).
- <sup>58</sup>J. K. Weltman and R. M. Dowben, *Proc. Natn. Acad. Sci., U.S.A.* **70**, 3230 (1973).
- <sup>59</sup>R. G. Adjodi, *Indian J. Exp. Biol.* **6**, 144 (1968).
- <sup>60</sup>M. S. Kerr, *Develop. Biol.* **20**, 1 (1969).
- <sup>61</sup>Y. Croisille, *Amer. Zool.* **14**, 1219 (1974).
- <sup>62</sup>H. J. Ceccaldi, *Rec. Trav. Stat. mar. Endoume*, **44,60**, 403 (1968).
- <sup>63</sup>H. W. Beams and R. G. Kessel, *J. Cell Biol.* **18**, 621 (1963).
- <sup>64</sup>F. J. Finamore and J. S. Clegg, *The Cell Cycle* (editors G. L. Padilla, G. L. Whitson and I. Cameron), Chap. 12. Academic Press, New York (1969).
- <sup>65</sup>A. H. Warner and D. K. McClean, *Develop. Biol.* **18**, 279 (1968).
- <sup>66</sup>W. Wieser, *J. Marine Biol. Assoc. UK* **45**, 507 (1965).
- <sup>67</sup>W. L. Lee, *Ecology* **47**, 930 (1966).
- <sup>68</sup>G. C. Genofre, *Compt. Rend Acad. Sci., Ser. D*, **276**, 2269 (1973).
- <sup>69</sup>B. H. Davies, W.-J. Hsu and C. O. Chichester, *Comp. Biochem. Physiol.* **33**, 601 (1970).
- <sup>70</sup>D. Arens, W. Seilmeier, F. Weber, G. Kloos and W. Grosch, *Biochim. Biophys. Acta* **327**, 295 (1973).
- <sup>71</sup>P. F. Zagalsky, H. J. Ceccaldi and R. Dumas, *Comp. Biochem. Physiol.* **34**, 579 (1970).
- <sup>72</sup>P. F. Zagalsky and P. J. Herring, *Comp. Biochem. Physiol.* **41B**, 397 (1972).
- <sup>73</sup>H. Nakagawa, M. Kayama and S. Askawa, *J. Fac. Fisheries Animal Husbandry, Hiroshima Univ.* **12**, 21 (1973).
- <sup>74</sup>H. Nakagawa, M. Kayama and S. Asakawa, *J. Fac. Fisheries Animal Husbandry, Hiroshima Univ.* **13**, 1 (1973).
- <sup>75</sup>H. Nakagawa, M. Kayama and S. Asakawa, *J. Fac. Fisheries Animal Husbandry, Hiroshima Univ.* **10**, 61 (1971).
- <sup>76</sup>H. J. Ceccaldi and P. F. Zagalsky, *Comp. Biochem. Physiol.* **21**, 435 (1967).
- <sup>77</sup>R. Kuhn and H. Kühn, *Angew. Chem.* **78**, 979 (1966).
- <sup>78</sup>R. Kuhn and H. Kühn, *Europ. J. Biochem.* **2**, 349 (1967).
- <sup>79</sup>M. Buchwald and W. P. Jencks, *Biochemistry* **7**, 844 (1968).
- <sup>80</sup>D. F. Cheesman, P. F. Zagalsky and H. J. Ceccaldi, *Proc. Roy. Soc. B*, **164**, 130 (1966).
- <sup>81</sup>W. P. Jencks and B. Buten, *Arch. Biochem. Biophys.* **107**, 511 (1964).
- <sup>82</sup>M. Buchwald and W. P. Jencks, *Biochemistry* **7**, 834 (1968).
- <sup>83</sup>R. M. Quarmby, Ph.D. Thesis, London University (1971).
- <sup>84</sup>R. M. Quarmby, D. Nordan, P. F. Zagalsky, H. J. Ceccaldi and R. Dumas, Paper in preparation.
- <sup>85</sup>T. Yoshimura and T. Isemura, *J. Biochem.* **69**, 969 (1971).
- <sup>86</sup>G. D. Sabato and N. O. Kaplan, *J. Biol. Chem.* **239**, 438 (1964).
- <sup>87</sup>J. Heller and J. Horwitz, *J. Biol. Chem.* **248**, 6308 (1973).
- <sup>88</sup>E. W. Abrahamson and S. E. Ostroy, *Prog. Biophys. Mol. Biol.* **17**, 181 (1967).
- <sup>89</sup>D. B. Gammack, J. H. Raper, P. F. Zagalsky and R. Quarmby, *Comp. Biochem. Physiol.* **40B**, 295 (1971).
- <sup>90</sup>A. Kropf, *Vision Res.* **7**, 811 (1967).
- <sup>91</sup>W. L. Lee and P. F. Zagalsky, *Biochem. J.* **101**, 9c (1966).
- <sup>92</sup>B. C. L. Weedon, *Carotenoids* (editor O. Isler), Chap. 5, Birkhäuser, Basel (1971).
- <sup>93</sup>M. H. Smith, *J. Theoret. Biol.* **13**, 261 (1966).
- <sup>94</sup>R. A. Capaldi and G. Vanderkooi, *Proc. Natn. Acad. Sci. U.S.A.* **69**, 930 (1972).
- <sup>95</sup>F. T. Hatch and A. L. Bruce, *Nature, Lond.* **218**, 1166 (1968).
- <sup>96</sup>C. G. Bigelow, *J. Theoret. Biol.* **16**, 187 (1967).
- <sup>97</sup>Y. Nozaki and C. Tanford, *J. Biol. Chem.* **246**, 2211 (1971).
- <sup>98</sup>A. Kropf, B. P. Whittenberger, S. P. Goff and A. S. Waggoner, *Exp. Eye Res.* **17**, 591 (1973).
- <sup>99</sup>C. Enzell and G. Francis, *Acta Chem. Scand.* **22**, 344 (1968).
- <sup>100</sup>P. Blatz, M. Lin, P. Balasubramanian, V. Balasubramanian and P. B. Dewhurst, *J. Am. chem. Soc.* **91**, 5930 (1969).

- <sup>101</sup>J. Horwitz and J. Heller, *J. Biol. Chem.* **249**, 4712 (1974).
- <sup>102</sup>P. J. Herring, *Comp. Biochem. Physiol.* **39B**, 1039 (1971).
- <sup>103</sup>B. A. Helm and P. F. Zagalsky, Unpublished observations.
- <sup>104</sup>A. G. Andrews, G. Borch, S. L. Jensen and G. Snatzke, *Acta Chem. Scand.* **B28**, 730 (1974).
- <sup>105</sup>M. Kanai, A. Raz and D. S. Goodman, *J. Clin. Invest.* **47**, 2025 (1968).
- <sup>106</sup>P. A. Peterson and I. Beggard, *J. Biol. Chem.* **246**, 25 (1971).
- <sup>107</sup>S. Futterman and J. Heller, *J. Biol. Chem.* **247**, 5168 (1972).
- <sup>108</sup>P. A. Peterson and L. Rask, *J. Biol. Chem.* **246**, 7544 (1971).
- <sup>109</sup>D. Klinger and M. Karplus, as in ref. 110.
- <sup>110</sup>B. Honig and T. G. Ebrey, *Ann. Rev. Biophys. Bioengr.* **3**, 151 (1974).
- <sup>111</sup>M. Takezaki and Y. Kito, *Nature, Lond.* **215**, 1197 (1967).
- <sup>112</sup>M. Azuma, K. Azuma and Y. Kito, *Biochim. Biophys. Acta* **351**, 133 (1974).
- <sup>113</sup>E. M. Johnson and R. Zand, *Biochemistry* **12**, 4637 (1973).
- <sup>114</sup>E. M. Johnson and R. Zand, *Biochemistry* **12**, 4631 (1973).
- <sup>115</sup>M. C. Gardiner and A. J. Thomson, Unpublished observations.
- <sup>116</sup>C. A. Coulson and D. K. Davies, *Trans. Faraday Soc.* **48**, 777 (1952).
- <sup>117</sup>P. Y. Chou and G. D. Fasman, *J. Mol. Biol.* **74**, 263 (1973).
- <sup>118</sup>P. N. Lewis and H. A. Scheraga, *Arch. Biochem. Biophys.* **144**, 576 (1971).
- <sup>119</sup>P. N. Lewis and H. A. Scheraga, *Biochim. Biophys. Acta.* **336**, 153 (1974).
- <sup>120</sup>P. N. Lewis, F. A. Momany and H. A. Scheraga, *Proc. Natn. Acad. Sci., U.S.A.* **68**, 2293 (1971).
- <sup>121</sup>K. Nagano, *J. Mol. Biol.* **75**, 401 (1973).
- <sup>122</sup>G. Némethy and H. A. Scheraga, *J. Phys. Chem.* **66**, 1773 (1962).
- <sup>123</sup>J. C. J. Bart and C. M. MacGillivray, *Acta Cryst.* **B24**, 1587 (1968).
- <sup>124</sup>S. J. Leach, G. Némethy and H. A. Scheraga, *Biopolymers* **4**, 887 (1966).
- <sup>125</sup>L. P. Vernon, B. Ke, H. H. Mollenhauer and E. R. Shaw, *Prog. Photosynthetic Res.* **1**, 137 (1969).
- <sup>126</sup>J. P. Thornber, J. C. Stewart, M. W. C. Hatton and J. Leggett-Bailey, *Biochemistry* **6**, 2006 (1967).
- <sup>127</sup>D. J. Haidak, C. K. Mathews and B. M. Sweeney, *Science* **152**, 212 (1966).
- <sup>128</sup>F. T. Haxo, J. H. Kycia, H. W. Siegelman and G. F. Somers, *Abstr. 3rd Intern. Symp. Carotenoids, Clug.* p. 73 (1972).
- <sup>129</sup>H. G. Garewal and A. R. Wasserman, *Biochemistry* **13**, 4072 (1974).
- <sup>130</sup>J. O. Erickson and P. E. Blatz, *Vision Res.* **8**, 1367 (1968).
- <sup>131</sup>B. Ke, *Meth. Enzym.* **23**, 624 (1971).
- <sup>132</sup>C. Subbarayan and H. R. Cama, *Indian J. Biochem.* **3**, 225 (1966).
- <sup>133</sup>T. H. Ji, J. L. Hess and A. A. Benson, *Biochim. Biophys. Acta* **150**, 676 (1968).
- <sup>134</sup>F. A. McEvoy and W. S. Lynn, *J. Biol. Chem.* **248**, 4568 (1973).
- <sup>135</sup>A. Hager and H. Perz, *Planta* **93**, 314 (1970).
- <sup>136</sup>K. Shibata, *Biochim. Biophys. Acta* **22**, 398 (1956).
- <sup>137</sup>A. Hager, *Planta* **91**, 38 (1970).
- <sup>138</sup>R. L. Hall, M. C. Kung, M. Fu, B. J. Hales and P. A. Loaches, *Photochem. Photobiol.* **18**, 505 (1973).
- <sup>139</sup>U. Schwenker, M. St-Onge and G. Gingras, *Biochim. Biophys. Acta* **351**, 246 (1974).
- <sup>140</sup>U. Schwenker and G. Gingras, *Biochem. Biophys. Res. Commun.* **51**, 94 (1973).
- <sup>141</sup>D. Thirkell and M. I. S. Hunter, *J. Gen. Microbiol.* **58**, 289 (1969).
- <sup>142</sup>D. Thirkell and M. I. S. Hunter, *J. Gen. Microbiol.* **58**, 293 (1969).
- <sup>143</sup>D. Thirkell and M. I. S. Hunter, *J. Gen. Microbiol.* **62**, 125 (1970).
- <sup>144</sup>A. L. Tappel and C. J. Dillard, *Lipids* **3**, 221 (1967).
- <sup>145</sup>J. Glover, *Vitams Horm.* **31**, 1 (1973).
- <sup>146</sup>T. Hara and R. Hara, *Nature, Lond.* **242**, 39 (1973).
- <sup>147</sup>D. Oesterhelt, M. Meentzen and L. Schuhmann, *Europ. J. Biochem.* **40**, 453 (1973).
- <sup>148</sup>D. Bownds, *Nature, Lond.* **216**, 1178 (1967).
- <sup>149</sup>J. C. Saari, *J. Cell Biol.* **63**, 480 (1974).
- <sup>150</sup>L. Rask, *Europ. J. Biochem.* **44**, 1 (1974).
- <sup>151</sup>H. Schichi, M. S. Lewis, F. Irreverre and A. L. Stone, *J. Biol. Chem.* **249**, 529 (1969).
- <sup>152</sup>W. Stoeckenius and W. H. Kanau, *J. Cell Biol.* **38**, 337 (1968).
- <sup>153</sup>H. Muhilal and J. Glover, *Trans. Biochem. Soc.* **3(5)**, 744 (1965).
- <sup>154</sup>L. Mateu, A. Tardieu, V. Luzzati, L. Aggerbeck and A. M. Scanu, *J. Molec. Biol.* **70**, 105 (1972).
- <sup>155</sup>J. P. Segrest, R. L. Jackson, J. D. Morrisett and A. M. Gotto, *FEBS Lett.* **38**, 247 (1973).
- <sup>156</sup>B. M. Gilchrist, P. F. Zagalsky and J. P. Prebble, Unpublished observations.