

## $\alpha$ -Chiral allylboronates: reagents for asymmetric synthesis

Reinhard W. Hoffmann


Fachbereich Chemie der Philipps-Universität Marburg  
 Hans-Meerwein-Straße, D - 3550 Marburg a.d. Lahn

**Abstract** - Allylboronates as well as crotylboronates add to aldehydes forming homoallyl alcohols. On addition of chiral  $\alpha$ -heterosubstituted allyl(crotyl)boronates to aldehydes the chirality is transferred to the new stereocenter of the homoallyl alcohol. The asymmetric induction that can be effected with  $\alpha$ -chloro-E-crotylboronates is in the order of 20 - 30 : 1.  $\alpha$ -Methoxy-E-crotylboronates as well as  $\alpha$ -methyl-Z-crotylboronates turned out to be even more effective.

### INTRODUCTION

Among the carbon-carbon-bond forming reactions, the addition of organometallic reagents to aldehydes and ketones occupies a prominent position. To this class of reactions belongs the addition of allylboronates to aldehydes (ref. 1), a reaction that stands out by its high degree of chemoselectivity, since ketones, esters, amides, epoxides, nitriles, are not touched under the usual reaction conditions.


**Scheme 1**




Another desirable factor for a reaction generating new stereogenic centers is stereoselectivity. Assuming that the addition of the allylboronate to an aldehyde proceeds via a cyclic six-membered transition state we reasoned that the decreased Lewis-acidity of the boron in the allylboronates would correspond to a transition state late on the reaction coordinate and that the short boron oxygen bond involved should render these transition states quite compact, amenable to a high degree of stereocontrol. A chirally modified allylboronate should differentiate the two enantiotopic faces of the aldehyde and allow an enantioselective formation of the homoallyl alcohols 1. An obvious point for attachment of the chiral information is the glycol component of the boronate. Earlier studies (ref. 2) using a camphor derived chiral glycol were quite encouraging: Scheme (2).

More recently, significant improvements have been realized by others (ref. 3) using tartrate derived chiral auxiliaries: Scheme (3).

**Scheme 2**




**Scheme 3**



W.R. Roush, J.A.C.S 1985


**$\alpha$ -SUBSTITUTED ALLYL BORONATES**

However, we felt that an intrinsically better approach to stereocontrol would be to place the element of chirality right onto the core of the transition state (ref. 4), i.e. we became interested in  $\alpha$ -chiral allylboronates 2. These could on reaction with aldehydes transfer their chirality to the newly formed stereogenic center. There are two competing diastereomeric transition states for this process, the energy difference of which determines the level of the asymmetric induction. These transition states differ by the equatorial versus axial arrangement of the group X. There is the useful corollary that the transition state with an equatorial X leads to the homoallyl alcohol 4 with an E-double bond, whereas the transition state with an axial X leads to 3 with a Z-double bond. Hence, the level of asymmetric induction can already be assessed in the racemic series simply by evaluating the E/Z-ratio of the product homoallyl alcohol: Scheme (4).




A variety of  $\alpha$ -hetero-substituted allylboronates were prepared as shown in Scheme (5).

Surprisingly, the addition of the  $\alpha$ -bromoallylboronate to aldehydes showed a high preference for the formation of the homoallyl alcohol with a Z-double bond (ref. 5): Scheme (6).

**Scheme 6**


R = CH <sub>3</sub>	78%	93 : 7
= CH <sub>3</sub> -CH <sub>2</sub>	82%	94 : 6
= (CH <sub>3</sub> ) <sub>2</sub> CH	83%	96 : 6
= C <sub>6</sub> H <sub>5</sub>	80%	97 : 3

Since the E/Z-ratio in the product homoallyl alcohols is equal to the asymmetric induction that would originate from a chiral  $\alpha$ -substituted allylboronate, the selectivities recorded in scheme (7) demonstrate that high levels of chirality transfer can be realized using  $\alpha$ -chiral allylboronates.



**Scheme 7**




On reaction of **5** of ca. 95 % e.e. with prochiral aldehydes transfer of chirality proceeded to our full satisfaction. The relative and the absolute configuration of the adduct to acetaldehyde has been proven by conversion into the known methyl  $\alpha$ -methyl- $\beta$ -hydroxy-butyrates: Scheme (12).

**Scheme 12**

At first glance the fact that the products contain a vinyl chloride functionality could be considered as a disadvantage in view of the low reactivity of this function. On the other hand, as these vinyl chlorides have a stereohomogenous Z-double bond, there was the challenge to take advantage of just this particular situation. In consequence, we devised some re-functionalisation schemes in which the vinyl chloride function is the starting point for the generation of highly substituted dihydropyran and tetrahydrofuran rings, c.f. schemes (13) and (14). As a bonus, the creation of the additional stereogenic centers proceeded under high asymmetric induction as well.


**Scheme 13****Scheme 14**

Now let us focus again on the main theme, the reaction of the crotylboronates with chiral aldehydes. On addition to the aldehyde **6** either triade of stereocenters (ref. 12) could be formed with a good level of selectivity. Of course, high selectivity in the matched pair (ref. 9), in which the asymmetric induction of the substrate and the reagent cooperate is not difficult to obtain: Scheme (15).

**Scheme 15**


Reagent control of diastereoselectivity in the mismatched pair requires that the asymmetric induction from the reagent is high enough to override the asymmetric induction from the substrate aldehyde. An inspection of the  $\Delta\Delta G^\ddagger$  values involved shows, where the weak points are: Scheme (16).

Scheme 16




An asymmetric induction of our reagent of 97 : 3 corresponds to only a  $\Delta\Delta G^\ddagger$  of 2.0 kcal. This becomes insufficient if a strong asymmetric induction of a chiral aldehyde is to be overcome. The following example is indicative of such a situation: Scheme (17).

Scheme 17




And in another case the aldehyde remained the winner in the struggle for stereocontrol: Scheme (18).

Scheme 18




Thus, we were forced to develop other  $\alpha$ -heterosubstituted E-crotylboronates with even higher asymmetric induction. In pursuit of this goal we succeeded to realize the following transformation: Scheme (19).

Scheme 19




Scheme 20



However, this substitution was accompanied by a loss of 4 - 10 % e.e. by racemisation. Nevertheless we won a reagent which has an asymmetric induction of at least 95 % and hopefully higher. We conclude this from the NMR-spectra of the enol ethers obtained after addition of  $\bar{7}$  to aldehydes, which show the presence of solely the Z-configured product: Scheme (20).

Scheme 21




With the reagent **7** in hand we could finally attain a significantly higher level of reagent control of diastereoselectivity on addition to the refractory aldehydes studied before: Scheme (21).

One principal aspect of reagent control of diastereoselectivity should not be overlooked: These reactions, in which the asymmetric induction of the substrate aldehyde has to be overturned proceed more slowly than the reaction of a matched pair. The reaction of a mismatched pair is forced to use a higher energy transition state (cf. scheme (16)). This means that in order to get decent conversions long reaction times or reactions under 4 kbar pressure had to be applied. Moreover, all kinds of side reactions rear their ugly heads. The more we feel satisfied with the selectivities that could be realized in the mismatched cases.

Yet we ought to point out a potentially more economical and easier way: Instead of fighting uphill using reagent control of diastereoselectivity to form the product of a mismatched pair, e.g. stereotriade **D** in Scheme (22), it could perhaps be easier to generate another stereotriade **A**, if this happens to be the product of a matched pair. In this case the product would easily be generated diastereomerically pure, but it obviously has the undesired configuration at the hydroxyl bearing carbon. Since the configuration of such an alcohol can be inverted by a number of ways, this reasoning makes the development of methods for the diastereo- and enantioselective generation of syn- $\beta$ -methyl homoallyl alcohols even more urgent.

### $\alpha$ -CHIRAL Z-CROTYLBORONATES

The stereotriades **C** and **D** in scheme (22) have become available using the  $\alpha$ -chiral-E-crotylboronates as described above. In order to reach the other two stereotriades **A** and **B** as well,  $\alpha$ -chiral-Z-crotylboronates are required. First representatives were prepared in the racemic series. On addition to aldehydes the homoallyl alcohols **9**, this time with an E-double bond, were formed in preference.




This is a consequence of that fact that the axial position for X in the transition state (cf. scheme (21)) is now seriously crowded causing 1,3-allylic strain. Highest selectivity can, hence, be obtained with the methyl derivative **8**, the methyl group resending least on stereoelectronic and steric grounds to be in the equatorial position: Scheme (24).

**Scheme 24**




The pentenyl borate **10** could be obtained in a state of high diastereomeric (enantiomeric) purity making recourse to the method and the kind help of D.S. Matteson (ref. 13): Scheme (25).

**Scheme 25**




The degree of chirality transfer that can be obtained with this reagent **10** turned out to be rewarding indeed: Scheme (26).

**Scheme 26**




The relative and absolute configuration of the resulting homoallyl alcohols were established by the application of the reagent **10** to the synthesis of invictolide: Scheme (27).

**Scheme 27**




**Scheme 28**



In due course, the reagent **10** has been utilized in the assembly of the stereotriade at C-4 to C-6 in mycinolide V, the aglycon of one of the mycinamycin antibiotics. The synthesis of this macrolide has recently been completed (ref. 14): Scheme (28).

En route to the C-4/C-6 stereotriade of mycinolide the various crotylboronate reagents have been employed. The results in scheme (29) nicely illustrate the improvement in diastereoselectivity that was realized by changing from an achiral crotylboronate to a chirally modified crotylboronate to the present generation of  $\alpha$ -chiral crotylboronates.

Scheme 29



### Acknowledgement

I would like to thank my coworkers who developed these allylboronates to the point that they became useful reagents. These coworkers are: S. Dresely, K. Ditrich, T. Herold, B. Hildebrandt, G. Köster, W. Ladner, B. Landmann, J. Lanz, U. Schrott, H.-J. Zeiß and R. Stürmer. This work could not have been pushed forward without the constant support from the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie.

### REFERENCES

1. E. Favre, M. Gaudemar, *Compt. Rend. hebd. Acad. Sci. Ser. C.* **263**, 1543 (1966).
2. R.W. Hoffmann, T. Herold, *Chem. Ber.* **114**, 375 (1981).
3. R.W. Roush, A.E. Walts, L.K. Hoong, *J. Am. Chem. Soc.* **107**, 8186 (1985); N. Ikeda, I. Arai, H. Yamamoto, *J. Am. Chem. Soc.* **108**, 483 (1986); W.R. Roush, R. Halterman, *J. Am. Chem. Soc.* **108**, 297 (1986).
4. M.M. Midland, S.B. Preston, *J. Am. Chem. Soc.* **104**, 2330 (1982); V.J. Jephcote, A.J. Pratt, E.J. Thomas, *Chem. Commun.* **1984**, 800.
5. R.W. Hoffmann, B. Landmann, *Chem. Ber.* **119**, 1039 (1986).
6. S.L. Schreiber, K. Satake, *J. Am. Chem. Soc.* **105**, 6723 (1983); K.N. Houk, N.G. Rondan, Y.-D. Wu, J.T. Metz, M.N. Paddon-Row, *Tetrahedron* **40**, 2257 (1984); S.J. Danishefsky, E. Larson, J.P. Springer, *J. Am. Chem. Soc.* **107**, 1274 (1985).
7. R.W. Hoffmann, *Angew. Chem.* **94**, 569 (1982); *Angew. Chem., Int. Ed. Engl.* **21**, 555 (1982).
8. R.W. Hoffmann, H.-J. Zeiß, *J. Org. Chem.* **46**, 1309 (1981).
9. S. Masamune, W. Choy, J.S. Petersen, L.R. Sita, *Angew. Chem.* **97**, 1 (1985); *Angew. Chem., Int. Ed. Engl.* **24**, 1 (1985).
10. D.J. Tsai, D.S. Matteson, *Organometallics* **2**, 236 (1983).
11. R.W. Hoffmann, S. Dresely, *Angew. Chem.* **98**, 186 (1986); *Angew. Chem., Int. Ed. Engl.* **25**, 189 (1986).
12. R.W. Hoffmann, *Angew. Chem.* **99**, in print (1987).
13. K.M. Sadhu, D.S. Matteson, G.D. Hurst, J.M. Kurosky, *Organometallics* **3**, 804 (1984).
14. K. Ditrich, T. Bube, R. Stürmer, R.W. Hoffmann, *Angew. Chem.* **98**, 1016 (1986); *Angew. Chem., Int. Ed. Engl.* **25**, 1028 (1986).